

Pre-Implantation Genetic Testing

Policy Number: AHS – M2039 – Pre-Implantation Genetic Testing	Policy Revision Date: 07/01/2025 Initial Policy Effective Date: 12/01/2024
---	---

[POLICY DESCRIPTION](#) | [RELATED POLICIES](#) | [INDICATIONS AND/OR LIMITATIONS OF COVERAGE](#) | [TABLE OF TERMINOLOGY](#) | [SCIENTIFIC BACKGROUND](#) | [GUIDELINES AND RECOMMENDATIONS](#) | [APPLICABLE STATE AND FEDERAL REGULATIONS](#) | [APPLICABLE CPT/HCPCS PROCEDURE CODES](#) | [EVIDENCE-BASED SCIENTIFIC REFERENCES](#) | [REVISION HISTORY](#)

I. Policy Description

Preimplantation genetic testing (PGT) involves the biopsy of a single cell, or a few cells of embryos to facilitate genetic testing for various genetic conditions. These conditions range from aneuploidies to monogenic disorders to structural deformities of the chromosomes themselves.¹

II. Related Policies

Policy Number	Policy Title
AHS-M2017	Genetic Testing for Cystic Fibrosis
AHS-M2145	General Genetic Testing, Germline Disorders
AHS-M2146	General Genetic Testing, Somatic Disorders
AHS-M2179	Prenatal Screening (Genetic)

III. Indications and/or Limitations of Coverage

Application of coverage criteria is dependent upon an individual’s benefit coverage at the time of the request. Specifications pertaining to Medicare and Medicaid can be found in the “Applicable State and Federal Regulations” section of this policy document.

- 1) Genetic counseling **IS REQUIRED** for individuals contemplating preimplantation genetic testing.
- 2) Preimplantation genetic testing for specific mutation(s) or chromosomal changes that have been associated with a specific disorder **MEETS COVERAGE CRITERIA** when one of the following conditions is met:
 - a) Both biological parents are known carriers of an early-onset, autosomal recessive disorder.
 - b) One biological parent is a known carrier of an early-onset, autosomal recessive disorder and the other biological parent is unavailable for testing.
 - c) One biological parent is a known carrier of an early-onset, autosomal recessive disorder and together, the biological parents have produced previous offspring affected with the disorder.
 - d) One biological parent is a known carrier of an early-onset, autosomal dominant disorder.
 - e) One biological parent is a known carrier of an early-onset, X-linked disorder.
 - f) One biological parent carries a balanced or unbalanced chromosomal translocation.

- g) One biological parent is a known carrier of one of the following adult-onset, autosomal dominant disorders:
 - i) Huntington Disease (*HTT*).
 - ii) Autosomal dominant polycystic kidney disease (*PKD, PKD2*).
- 3) In the absence of an early-onset, sex-linked disorder, preimplantation genetic testing for sex selection **DOES NOT MEET COVERAGE CRITERIA**.
- 4) Preimplantation genetic testing **DOES NOT MEET COVERAGE CRITERIA** for any of the following situations:
 - a) Preimplantation genetic testing for adult-onset disorders not discussed above.
 - b) Preimplantation HLA genotyping for purposes of identifying potential tissue or organ donors.
 - c) Routine preimplantation screening for chromosomal abnormalities, including testing based on advanced maternal age.

The following does not meet coverage criteria due to a lack of available published scientific literature confirming that the test(s) is/are required and beneficial for the diagnosis and treatment of an individual's illness.

- 5) For all other situations not described above, preimplantation genetic testing **DOES NOT MEET COVERAGE CRITERIA**.

IV. Table of Terminology

Term	Definition
aCGH	Array-based comparative genomic hybridization
ACMG	American College of Medical Genetics and Genomics
AD	Autosomal dominant
AR	Autosomal recessive
ART	Assisted reproductive technology
ASRM	American Society of Reproductive Medicine
BFS	British Fertility Society
BMI	Body mass index
cLBR	Cumulative live birth rate
CLIA '88	Clinical Laboratory Improvement Amendments of 1988
CMS	Centers for Medicare and Medicaid
DNA	Deoxyribose nucleic acid
ESHRE	European Society of Human Reproduction and Embryology
HLA	Human leukocyte antigen
ISAR	Indian Society for Assisted Reproduction
IVF	In-vitro-fertilization

LDTs	Laboratory-developed tests
NGS	Next-generation sequencing
PCR	Polymerase chain reaction
PGD	Preimplantation genetic diagnosis
PGD-A	Preimplantation genetic diagnosis for aneuploidy testing
PGDIS	Preimplantation Genetic Diagnosis International Society
PGS	Preimplantation genetic screening
PGT	Preimplantation genetic testing
PGT-A	Preimplantation genetic test for aneuploidy
PGT-M	Preimplantation genetic testing for monogenic defects
PGT-SR	Preimplantation genetic testing-structural rearrangements
qPCR	Quantitative real-time PCR
RPL	Recurrent pregnancy loss
SNP	Single nucleotide polymorphism
CCMG	Canadian College of Medical Geneticists
XL	X-linked

V. Scientific Background

Preimplantation genetic testing (PGT) in conjunction with assisted reproductive technology (ART) was developed to allow reproductive partners at-risk of transmitting a genetic condition to their offspring to have an unaffected child without facing prenatal diagnosis and termination of pregnancy.² Initially offered for diagnosis in couples at-risk for single gene genetic disorders, such as cystic fibrosis, spinal muscular atrophy and Huntington disease, preimplantation genetic diagnosis (PGD) has most frequently been employed in assisted reproduction as preimplantation genetic screening (PGS) for detection of chromosome aneuploidy from advancing maternal age or structural chromosome rearrangements. The Preimplantation Genetic Diagnosis International Society (PGDIS) estimates that nearly 80% of PGT cycles have been performed for aneuploidy screening, 12% for single gene disorders, 6% for chromosome rearrangements and 2% for sibling human leukocyte antigen (HLA) matching.² Both this and the European Society of Human Reproduction and Embryology (ESHRE) surveys confirm that aneuploidy testing is the major indication for PGT.³

Embryonic genetic material used for PGT can be obtained from any of three sources: polar bodies from oocytes, blastomeres from day two or three, or trophoctoderm cells from blastocysts.⁴ Polar bodies are typically analyzed if the embryo cannot be biopsied. However, polar body analysis is only useful for finding maternally inherited mutations or a cell division error during oocyte development. Furthermore, since genetic changes occur after the polar body develops, test results are of limited use. Additionally, as many as 30% of oocytes will not fertilize successfully, causing the test to fail.¹

Blastomeres from day two or three (cleavage stage) were once the preferred practice in in-vitro-fertilization (IVF) as more embryos survived in culture by day three compared to days five or six (blastocyst stage). Despite the greater survival rate of day three embryos, these embryos were found to have a lower survival rate in a sustained implantation compared to day five embryos. Overall, trophoctoderm biopsy on day five is preferable as it has no measurable impact on embryo

development.⁵ Up to two or three dozen cells can be removed without disrupting development; although, common practice is to remove five to eight cells. Day five and later embryos also provide more DNA for testing compared to other stages of development.¹ Improved results have been seen with decreasing use of day three blastomere biopsy in favor of day five trophectoderm biopsy.⁴

PGS is emerging as one of the most valuable tools to enhance pregnancy success with assisted reproductive technologies by assessing embryos for aneuploidy.⁶

As the genetic basis of more disorders are identified, increasing demand for and acceptance of the use of PGT for adult-onset disorders, such as Huntington disease, hereditary breast and ovarian cancer and Alzheimer disease, have occurred. Using PGD to screen embryos for diseases or mutations that confer an increased risk for developing a particular disease raises issues of how to weigh the benefits of PGD to the future child against the risks of PGD and ART.³ The Ethics committee for the American Society of Reproductive Medicine (ASRM) found that PGD for adult-onset conditions is ethically justifiable when the conditions are serious and when there are no known interventions for the conditions or the available interventions are either inadequately effective or significantly burdensome.⁷ The use of PGT for nonmedical sex selection or family balancing continues to be controversial, and the Ethics committee has stated that it is acceptable for facilities to offer this service; however, employees wishing to decline participation in these procedures should be allowed to do so.⁸

Women recommended for preimplantation genetic diagnosis for aneuploidy testing (PGD-A) or PGT for aneuploidy (PGT-A) are those of advanced reproductive age with a history of recurrent miscarriages and/or IVF failures; PGD-A is currently performed on trophectoderm biopsies by 24 different chromosome screening techniques.⁹ Trophectoderm biopsies are a safe and extensively validated approach with a low margin of error and miscarriage rate as well as a suspected high sustained pregnancy rate.⁹ However, Alteri, et al. (2019) states that while PGT-A allows for an increased implantation rate, current data does not show an increase in successful pregnancy rates. Researchers agree that this technology is imperfect as Ledger (2019) reports that PGD-A can incorrectly designate an euploid embryo as an aneuploid embryo, leading to the unnecessary waste of embryos. These researchers also suggest that this type of screening may only be necessary in women between the ages of 35 and 44, as embryonic aneuploidy rates are low below 37 years of age and “costly screening for aneuploid seems pointless for women over 44 years of age, as almost all embryos are aneuploid.”¹¹

The development of whole genome amplification and genomic tools, such as single nucleotide polymorphism (SNP) microarrays and comparative genomic hybridization microarrays, has led to faster, more accurate diagnoses that lead to improved pregnancy and live birth rates.¹² Next-generation sequencing has also been used to distinguish between normal and abnormal embryos.¹³ PGD-polymerase chain reaction (PCR) is often used to amplify the obtained DNA from the blastomere biopsy for further analysis.¹⁴ Fluorescence In-situ Hybridization (FISH) has also been used for PGD and is an efficient method that may help to decrease IVF failure in infertile patients.¹⁵

Other researchers are attempting to develop a non-invasive preimplantation genetic testing technique. Farra, et al. (2018) state that circulating cell-free embryonic DNA can be obtained from used culture media from blastocysts and in blastocoel fluid; this can then be used as a non-invasive method to evaluate genetic embryonic properties.

Proprietary Testing

Companies, such as Natera, have developed a preimplantation genetic test for monogenic/single gene conditions called Spectrum™ that includes PGT-A, PGT-M, and PGT-SR. This PGT-A uses SNP microarray technology.¹⁷ Simon, et al. (2018) studied IVF outcomes with this test when measuring PGT-A and euploid embryo transfer in day five or six embryos. An implantation rate and live birth rate of 69.9% and 64.5%, respectively, was identified.¹⁸ The authors concluded that “SNP-based PGT-A can mitigate the negative effects of maternal age on IVF outcomes in cycles with transfer, and that pregnancy outcomes from SET [single-embryo transfer] cycles are not significantly different from those of double-embryo transfer cycles, and support the use of SET when transfers are combined with SNP-based PGT-A.”¹⁸

iGLS Reproductive Genetics developed a PGT-A that uses next-generation sequencing (NGS) to analyze thousands of DNA sequences that are unique to each chromosome allowing for the accurate identification of extra or missing chromosomes.¹⁹

PacGenomics has developed several PGT's for aneuploidies (PGT-A), structural chromosomal rearrangements (PGT-SR) with a subsequent PGT-SR Plus® which is used to differentiate between normal and balanced translocation carrier embryos in translocation cases, and for monogenic/single gene disorders (PGT-M). PGT-A for aneuploidies is performed on embryos created through IVF to screen for chromosomal abnormalities. PGT-SR is a genetic test performed on embryos created through IVF to screen for chromosomal structural rearrangements caused by balanced translocations and inversions. PGT-SR-Plus® can be added to PGT-SR which helps identify translocation carriers in patients who are not initially suspected to be at significant risk. PGT-M is performed on embryos created through IVF that is designed for individuals who know they are at an increased risk of having a child with a specific genetic disorder.²⁰

Reproductive Genetic Innovations (RGI) developed several PGT tests including PGT-A, PGT-SR, and PGT-M which has similar functions to the other proprietary tests. RGI provides an additional test, PGT-HLA, which identifies embryos that are HLA compatible with a child who needs a bone marrow or cord blood transplant to help treat blood disorders.²¹

Clinical Utility and Validity

Dreesen, et al. (2014) performed a study assessing the accuracy of diagnoses made based on PGD. A total of 940 cases covering 53 genetic disorders were re-evaluated using a PCR-based test. Of the 940 embryos, 881 (93.7%) of these embryos had two agreeing diagnoses. The first evaluation breakdown was 234 unaffected embryos, 590 affected, and 116 aberrant whereas the re-evaluation's breakdown was 283 unaffected embryos, 578 affected, and 79 aberrant. The sensitivity of this method was 99.2%, and its specificity was 80.2%. Allelic drop-out, mosaicism, and human error were the three most common causes of error.²²

Ghiossi, et al. (2018) performed a study focusing on couples' decisions based on expanded carrier screening. Forty-five couples took a survey of their reproductive decision making after receiving their results, and of those 45, 28 said they would plan IVF with PGD or a prenatal diagnosis in future pregnancies. Of the 19 pregnant respondents, eight chose a prenatal diagnosis route, two planned amniocenteses but miscarried, and nine considered the condition insufficiently severe to warrant invasive testing. Three of the eight that chose the prenatal diagnosis route were affected by a condition, and two pregnancies were terminated. Disease severity was found to be a significant association with

changes in decision making. Thirteen respondents did not plan to use the results from the carrier screening and four responses were unclear.²³

Kamath, et al. (2019) analyzed 207,697 data sets from women undergoing single-embryo transfer after PGT or IVF without PGT between the year 2000 and 2016. Results showed a significantly higher incidence of zygotic splitting following PGT (2.4%) compared to following non-PGT IVF (1.5%); this shows “a likely increased risk of monozygotic splitting following embryo biopsy”²⁴ and highlights a potential risk with PGT embryonic biopsies.

A new study was published which compared the live birth rates of embryos fertilized without chromosome analysis compared to those analyzed via PGT-A and comprehensive chromosome screening of the first and second polar body.²⁵ All mothers were of advanced maternal age between 36 and 40 years old. A total of 396 women enrolled in this multicenter, randomized clinical trial. Two hundred and five women had chromosomal screening, and 50 (24%) had a live birth within a year; in the group without intervention, which was comprised of 191 women, 45 (24%) had a live birth within a year.²⁵ It is important to note that the groups had a slightly different number of participants. This study shows that PGT-A allows for similar birth rates when compared to embryos fertilized without chromosome analysis via intracytoplasmic sperm injection (ICSI). However, the authors warn that “Whether these benefits outweigh drawbacks such as the cost for the patient, the higher workload for the IVF lab and the potential effect on the children born after prolonged culture and/or cryopreservation remains to be shown.”²⁵

A meta-analysis focusing on evaluating the effectiveness and safety of PGT-A in women undergoing an IVF treatment was conducted in 2020. Thirteen randomized controlled trials—involving a total of 2794 women—reporting data on clinical outcomes were included. The meta-analysis concluded that there existed insufficient evidence for preimplantation genetic testing for abnormal chromosomes numbers to provide a difference in cumulative live birth rate, live birth rate after the first embryo transfer, or miscarriage rate between IVF with and IVF without PGT-A as currently performed, and therefore “the effect of PGT-A on clinical pregnancy rate is uncertain.” The evidence evinced that though the observed cumulative live birth rate (cLBR) was 24% in the control group, the chance of live birth following the results of one IVF cycle with PGT-A is between 17% and 34%. Similarly, trials focusing on IVF with addition of PGT-A boasted an average cLBR of 29% in the control group, but the chance of live birth following the results of one IVF cycle with PGT-A was between 12% and 29%. When PGT-A is performed with FISH, the chance of live births after the first transfer in the control group (31%) fell to between 16% and 29% for those tested. Thus, the authors caution that “Women need to be aware that it is uncertain whether PGT-A with the use of genome-wide analyses is an effective addition to IVF, especially in view of the invasiveness and costs involved in PGT-A”, going so far as to state that “PGT-A using FISH for the genetic analysis is probably harmful.”²⁶

NGS can be used for PGS to screen for aneuploidies in IVF scenarios. A study by Yap, et al. (2019) analyzed results from a total of 391 IVF pregnancies whose embryos were cultured to the blastocyst stage; a total of 1361 blastocysts were analyzed.²⁷ Of the 1361 blastocysts, 423 were identified as aneuploid, 723 as euploid and 216 as mosaic (contained varying cell lines).²⁷ These results show that NGS can be used to identify mosaic and aneuploid blastocysts and is an effective PGS tool.

Zeevi, et al. (2021) studied the clinical validity of Haploseek, a method for preimplantation genetic testing and compared the results to PCR-based PGT case results. A total of 151 embryo biopsies from 27 PGT cases were obtained and sequenced using Haploseek to predict the chromosome copy number

variants (CNVs) and relevant variant-flanking haplotypes in each embryo. For each of the 151 embryo biopsies, all Haploseek-derived haplotypes and CNVs were concordant with clinical PGT results. The authors conclude that “Haploseek is clinically accurate and fit for all standard clinical PGT applications.”²⁸

Kumar, et al. (2022) noted the lack of comprehensive embryo genetic assessment in PGT. The authors used parental genome sequencing and embryo genotyping to create a whole genome reconstruction. Using a combination of molecular and statistic techniques, the authors could infer inherited genome sequences and model susceptibility to common conditions. The study included 110 embryos from ten couples and investigated 12 common conditions including cancer and autoimmune diseases. The method resulted in “genotype accuracy of 99.0–99.4% at sites relevant to polygenic risk scoring in cases from day-5 embryo biopsies and 97.2–99.1% in cases from day-3 embryo biopsies.” The authors conclude that these results can “inform the discussion of utility and implementation of genome-based PGT in clinical practice.”²⁹

VI. Guidelines and Recommendations

American College of Obstetricians and Gynecologists (ACOG)

In 2017, the ACOG noted that if a carrier couple (carriers for the same condition) is identified, genetic counseling is encouraged so that options such as PGD or prenatal diagnosis may be discussed. This guideline was reaffirmed in 2023.³⁰

In 2020, the ACOG published a series of recommendations in their “ACOG Committee Opinion” Number 799. The committee opinion was reaffirmed in 2023. These recommendations are shortened for brevity and reported below:

- “Preimplantation genetic testing comprises a group of genetic assays used to evaluate embryos before transfer to the uterus. Preimplantation genetic testing-monogenic (known as PGT-M) is targeted to single gene disorders. Preimplantation genetic testing-monogenic uses only a few cells from the early embryo, usually at the blastocyst stage, and misdiagnosis is possible but rare with modern techniques. Confirmation of preimplantation genetic testing-monogenic results with chorionic villus sampling (CVS) or amniocentesis should be offered.”
- “To detect structural chromosomal abnormalities such as translocations, preimplantation genetic testing-structural rearrangements (known as PGT-SR) is used. Confirmation of preimplantation genetic testing-structural rearrangements results with CVS or amniocentesis should be offered.”
- “The main purpose of preimplantation genetic testing-aneuploidy (known as PGT-A) is to screen embryos for whole chromosome abnormalities. Traditional diagnostic testing or screening for aneuploidy should be offered to all patients who have had preimplantation genetic testing-aneuploidy, in accordance with recommendations for all pregnant patients.”³¹

ACOG also notes the following: “Preimplantation genetic testing raises several ethical issues that are beyond the scope of this document. For example, there are complexities to offering preimplantation genetic testing to screen for adult-onset disorders or to determine transplantation compatibility for ill family members. Such issues are addressed extensively in the bioethics literature as well as documents from ACOG³² and the American Society of Reproductive Medicine.”^{31,33}

In 2008, ACOG published “Ethical Issues in Genetic Testing”, which was reaffirmed in 2020. In this guideline, ACOG published the following recommendations pertaining to prenatal testing for adult-onset disease. This guideline does not discuss ethics of preimplantation genetic testing:

- “Genetic testing of the fetus offers both opportunities and ethical challenges. Preconception and prenatal genetic screening and testing are recommended for a limited number of severe child-onset diseases because such screening and testing provides individuals with the chance to pursue assisted reproductive technology in order to avoid conception of an affected child, to consider termination of a pregnancy, or to prepare for the birth of a chronically ill child. With advancing genetic technology, however, physicians may increasingly face requests for testing of fetuses for less severe child-onset conditions, adult-onset conditions, or genetically linked traits.
- Principles regarding testing of children provide some guidance for when prenatal testing might be appropriate but this decision is significantly complicated by the various purposes that prenatal testing can have: to detect a fetal condition for pregnancy termination, to allow patients to prepare for the birth and care of a potentially affected child, or, more rarely, to detect and treat a fetal condition in utero. Furthermore, many times, a woman’s intentions regarding pregnancy termination evolve as genetic information becomes available to her. Therefore, testing the fetus for adult-onset disorders with no known therapeutic or preventive treatment (save prevention by pregnancy termination) should raise caution in a way similar to the manner in which testing of children can. In pregnancies likely to be carried to term, consideration should be given to whether, as in the case of testing children, the decision to test should be reserved for the child to make upon reaching adulthood. However, consideration also should be given to personal preference, that is, the interests individuals may have in terminating a pregnancy that may result in a life (such as life that will be affected by Huntington chorea) that they feel morally obliged or prefer not to bring into the world. Because these often are wrenching decisions for parents, referral to parent support networks (eg, National Down Syndrome Society, if that is the diagnosis of concern), counselors, social workers, or clergy may provide additional information and support.”³²

European Society for Human Reproduction and Embryology (ESHRE) PGD Consortium

In 2010, the ESHRE issued detailed guidelines related to technical aspects of PGD, specifically for the use of amplification techniques and for FISH. The ESHRE recommends that “misdiagnosis rates should be calculated for each type of assay and for all assays from a particular centre.” Additionally, they note that “Follow-up of pregnancies (including multiple pregnancy rate and outcome), deliveries, and the health of children at birth and beyond should be attempted and maintained along with the cycle data.”³⁴

In 2020, the ESHRE expanded upon their practice recommendations for preimplantation genetic testing. For the organization of PGT, the ESHRE provided patient inclusion/exclusion criteria. In general, “It is recommended that PGT is only applied when genetic diagnosis is technically feasible, and the reliability of the diagnosis is high. Current procedures in most IVF/PGT centres allow for overall error rates (resulting in misdiagnosis) as low as 1 to 3%. Each centre should be aware of their error rates and include this information in their informed consents and reports in an open communication with the patient.

When considering PGT, safety issues, female age, impossibility to retrieve male or female gametes, body mass index (BMI) and other contraindications for IVF should be considered as possible exclusion criteria.

Furthermore, exclusion from PGT should be considered if the woman has serious signs and symptoms of an autosomal dominant or X-linked disorder (for which PGT is requested), which could introduce medical complications during ovarian stimulation, oocyte retrieval or pregnancy or medical risks at birth. PGT should be carefully considered if one of the partners has serious physical or psychological problems, either linked to the tested disease or due to other conditions.”

Different preimplantation genetic testing for specific defects and disorders carried their own caveats and recommendations in terms of inclusion and exclusion of patients:

For *PGT-M, mitochondrial disorders and HLA*: “Cases of genetic variants of unknown significance that are not predictive of a phenotype should be excluded from PGT. PGT testing is inappropriate in case of uncertain genetic diagnosis (for example genetic/molecular heterogeneity), or in case of uncertain mode of inheritance.

For autosomal recessive disorders, where a single pathogenic variant has been diagnosed in the proband and only one parent, it is acceptable to offer PGT if the pathogenic genotype is attributed to a single gene and sufficient evidence from the family pedigree allows identification of the disease-associated haplotypes. Similarly, it is acceptable to offer PGT for known X-linked recessive single gene disorders with a clear unequivocal clinical diagnosis where no pathogenic variant was found in the proband but low- and high-risk haplotypes can be identified based on the family history.

Exclusion or non-disclosure testing can be indicated for late-onset disorders, such as Huntington’s disease, to avoid pre-symptomatic testing of the partner with a family history of the disease. Exclusion testing is preferred over PGT with non-disclosure of the direct test results to the couple.”

For *PGT for mitochondrial disorders*: “PGT is not indicated in case of homoplasmy. In cases where the causative pathogenic variant of the mitochondrial disease is encoded by nuclear DNA, testing is the same as for other monogenic disorders.”

For *HLA Typing*: “When all other clinical options have been exhausted, selection of HLA-matched embryos via PGT is acceptable for couples who already have a child affected with a malignant, acquired disorder or a genetic disorder where the affected child is likely to be cured or life expectancy is substantially prolonged by transplantation with stem cells from an HLA-matched sibling. Testing can be performed for HLA typing alone, if the recurrence risk of the disease is low, or in combination with autosomal dominant/recessive or X-linked disorders.”

For *PGT-SR*: “Depending on the technology used (FISH, quantitative real-time PCR (qPCR), comprehensive testing methods [array-based comparative genomic hybridisation (aCGH), single nucleotide polymorphism (SNP) array or next-generation sequencing (NGS)], different inclusion/exclusion criteria may apply. In general, PGT-SR is only recommended if the technique applied is able to detect all expected unbalanced forms of the chromosomal rearrangement. When comprehensive testing strategies are applied, it is acceptable to use information on copy number of nonindication chromosomes to refine embryo transfer strategies.”

For *PGT-A*: “For all, but in particular for RIF, RM and SMF couples, a previous karyotype of both partners is recommended since there is a higher chance of structural rearrangements for these indications. If an abnormal karyotype is identified, the technology for the detection of unbalanced abnormalities can differ from the regular PGT-A.”³⁵

Ethics Committee of the American Society for Reproductive Medicine (ASRM)

The ASRM Ethics committee has published several opinion guidelines over the years.

In 2018, the ASRM published a committee opinion on the use of preimplantation genetic testing for monogenic defects (PGT-M) for adult-onset conditions. These guidelines stated the following:

- “Preimplantation genetic testing for monogenic disease (PGT-M) for adult-onset conditions is ethically justified when the condition is serious and no safe, effective interventions are available.”
- “Reproductive liberty arguments ethically allow for PGT-M for adult-onset conditions of lesser severity or penetrance. In the latter cases, the application of the technology hinges on the evidence that PGT-M is a relatively low-risk procedure; this evidence may change.”
- “The Committee to strongly recommend that an experienced genetic counselor with knowledge about PGT-M play a major role in counseling patients considering such procedures.”³³

The ASRM also published an updated committee opinion in July 2023 detailing indications and management of preimplantation genetic testing for monogenic conditions, not limited to adult-onset. These recommendations state the following:

- “Suggested categorization of PGT-M indications (derived from table):
 - Childhood-onset, lethal or severe condition
 - Serious adult-onset conditions
 - Mild conditions or indications of limited or questionable risk reduction:
 - Low penetrance or susceptibility genes
 - Mild genetic variants
 - Carrier status for autosomal recessive conditions with carrier manifestations
 - Mild, common, and/or treatable conditions
 - Variants appearing to be de novo in an affected child
 - Variants of uncertain significance
- Indications for which PGT-M is not recommended:
 - Autosomal recessive carrier status without manifestations of symptoms
 - Combination of variants not associated with disease
 - Pseudodeficiency alleles
 - Somatic-only variants”
- “Preimplantation genetic testing for monogenic conditions should be offered if a significant reproductive risk is identified. Acceptance of PGT-M should be optional”

However, with these categorizations, the committee reiterates some key considerations surrounding the application of PGT-M and the ethical arguments in certain situations: “The initial application of PGT-M was primarily to prevent the transmission of severe, untreatable, or life-threatening childhood-onset conditions. Today, however, the use of the technology extends to a much broader range of genetic conditions for which support of PGT-M is less clear or even controversial, including those with mild to moderate phenotypes and later age of onset and of much greater clinical variability and/or reduced penetrance. A survey of laboratory genetic counselors in 2021 found that all participants believed that PGT-M should be allowed for conditions of lower penetrance, citing patient autonomy as a primary consideration. At this time, no guidelines exist to offer direction to those involved in determining when to offer or decline to perform PGT-M, although some ethical arguments exist. As such, physicians, genetic counselors, and PGT laboratories may develop internal policies

regarding PGT-M availability within their institution. If certain types of conditions or clinical scenarios are deemed not appropriate for PGT-M, the patient should be informed of the policy as early in the process as possible.”³⁶

Similar guidelines were also published in 2013 by the ASRM regarding the use of PGD for serious adult-onset conditions:

- “Preimplantation genetic diagnosis (PGD) for adult-onset conditions is ethically justifiable when the conditions are serious and when there are no known interventions for the conditions or the available interventions are either inadequately effective or significantly burdensome.
- For conditions that are less serious or of lower penetrance, PGD for adult-onset conditions is ethically acceptable as a matter of reproductive liberty. It should be discouraged, however, if the risks of PGD are found to be more than merely speculative.
- Physicians and patients should be aware that much remains unknown about the long-term effects of embryo biopsy on any developing fetus. Though thought to be without serious side effects, PGD for adult-onset diseases of variable penetrance should only be considered after patients are carefully and thoroughly counseled to weigh the risks of what is unknown about the technology and the biopsy itself against the expected benefit of its use.
- It is important to involve the participation of a genetic counselor experienced in such conditions before patients undertake PGD. Counseling should also address the patient specific prognosis for achieving pregnancy and birth through in-vitro-fertilization (IVF) with PGD.”³⁷

Additional guidelines were published in 2008, which stated the following:

- “Before PGD is performed, genetic counseling must be provided to ensure that patients fully understand the risk for having an affected child, the impact of the disease on an affected child, and the limitations of available options that may help to avoid the birth of an affected child.
- Prenatal diagnostic testing to confirm the results of PGD is encouraged strongly because the methods used for PGD have technical limitations that include the possibility for a false negative result.
- Available evidence does not support the use of PGS as currently performed to improve live birth rates in patients with advanced maternal age.
- Available evidence does not support the use of PGS as currently performed to improve live birth rates in patients with previous implantation failure.
- Available evidence does not support the use of PGS as currently performed to reduce miscarriage rates in patients with recurrent pregnancy loss related to aneuploidy.”³⁸

The ASRM Ethics Committee released a 2024 opinion on the use of preimplantation genetic testing for monogenic adult-onset conditions. The committee concluded that “PGT-M for adult-onset conditions that are most often fully penetrant or confer disease predisposition is ethically justified. Considerations including patient autonomy and the importance of supporting reproductive liberty dictate that decisions regarding the use of PGT-M in these situations should be made by patients as they consider the risk of disease development, the role of disease severity, and the age of onset.”³⁹

Concerning PGT-M, ASRM provided the following discussion points:

“Initially, PGT-M was developed to identify embryos resulting from in vitro fertilization (IVF) therapy cycles that carried genes for serious childhood-onset diseases. Preimplantation genetic testing for

monogenic diseases has been used recently for adult-onset single-gene disorders and disease predispositions. Examples include diseases that are not always but most often fully penetrant (e.g., Huntington disease and polycystic kidney disease) and genes that indicate a predisposition for cancer (e.g., BRCA1 and BRCA2 gene variants). Although Huntington disease is an autosomal dominant condition that is uniformly fatal, the age of onset varies with the pathogenic variant. In contrast, for cancer syndromes such as BRCA1 or BRCA2-associated hereditary breast and ovarian cancer, the presence of the identified gene variant(s) does not predict with certainty that an individual will ever develop the disease but significantly increases the lifetime risks for associated cancers. Moreover, some conditions can be treated successfully and may not be ultimately fatal or significantly affect the quality of life. For other conditions, successful treatments may be developed over time to prevent or treat the conditions for which the offspring are at risk. The use of PGT-M for serious adult-onset conditions thus raises challenging policy and ethical questions, given what is known about the human genome, disease etiology, the pace of medical progress, and embryo biopsy procedures.”³⁹

“Professional organizations such as the American Academy of Pediatrics currently recommend that genetic testing of children for adult-onset conditions for which interventions are unavailable is inappropriate until children reach 18 years of age and adulthood. This argument is based on the idea that the child has the right to an open future that is not burdened with the knowledge of a genetic condition. . . With some late-onset conditions, testing may be medically indicated before the child reaches adulthood. For example, autosomal dominant polycystic kidney disease (ADPKD) results in a progressive decline in kidney function, with kidney failure tending to occur in middle age. Children with ADPKD may not have any symptoms during childhood but require interval blood pressure assessment and renal ultrasonography beginning in childhood to monitor for early signs of kidney damage. . . Patients with a genetic condition often require repeated testing and treatment, often from early adulthood or beyond. For example, the financial and public health burdens of ADPKD include lost wages as well as long-term medical treatment. . . Individuals whose embryos are undergoing PGT-M for adult-onset diseases such as Huntington disease may also request that their own carrier status not be disclosed to them. It is ethically acceptable to honor such requests, but this practice remains controversial.”³⁹

“There are ethical reasons against using PGT-M for adult-onset conditions that are fully penetrant or confer disease predisposition. It is impossible to predict whether effective treatment modalities will be available before the manifestation of identified conditions by the time the offspring reach adulthood. Individuals with the disease-causing genetic variant may live healthy lives for several decades before a disease becomes an active concern in adulthood. Moreover, some of these genes may have variable expressivity or reduced penetrance, manifesting as a much milder form of illness than anticipated or perhaps never expressing illness at all, as in the case of some pathogenic variants that increase the lifetime risk of cancer. Genes that indicate a predisposition to cancer, such as *BRCA1* and *BRCA2*, present unique challenges. The current understanding of the complex interactions between DNA and the environment is limited. A woman who carries a *BRCA1* or *BRCA2* pathogenic variant has an increased lifetime risk of developing breast and/or ovarian cancer but may never develop these cancers for reasons that are not yet fully understood. . . Critics of PGT-M also argue that using the procedure for embryo selection risks devaluing certain lives. They contend that PGT-M can potentially send a negative message regarding the value of those individuals living with the disease, including those who have the mutation for the disease but have not yet developed a physical manifestation of the syndrome.”³⁹

Additionally, the Committee recommends “an experienced preimplantation genetic testing genetic counselor with knowledge about both the condition and assisted reproductive technology treatment

play a significant role in the counseling of prospective patients considering using PGT-M for adult-onset conditions. Counseling from medical professionals with expertise in the condition should also be considered as appropriate.”³⁹

Preimplantation Genetic Diagnosis International Society (PGDIS)

In the report from the 2021 PGDIS Expert Consultation on Mosaic Embryo Transfer, the PGDIS published a position statement on the transfer of mosaic embryos, including the following recommendations for clinicians:

- “Patients should continue to be advised that any genetic test based on sampling one or small number of cells biopsied from preimplantation embryos cannot be 100% accurate because of a combination of technical and biological factors, including cell mosaicism;
- Patient information and consent forms for aneuploidy testing should be modified to include the possibility of mosaic results; and
- In general, transfer of blastocysts with a normal euploid result should be prioritized over those with mosaic results unless other indications, such as patient preference, are raised.”⁴⁰

American College of Medical Genetics and Genomics (ACMG)

The ACMG has released guidelines on prenatal/preconception carrier screening, primarily when to test:

- “Disorders should be of a nature that most at-risk patients and their partners identified in the screening program would consider having a prenatal diagnosis to facilitate making decisions surrounding reproduction.
- For each disorder, the causative gene(s), mutations, and mutation frequencies should be known in the population being tested, so that meaningful residual risk in individuals who test negative can be assessed.”
- There must be validated clinical association between the mutation(s) detected and the severity of the disorder.”⁴¹

In 2021 the ACMG published recommendations on screening for autosomal recessive and X-linked conditions during pregnancy and preconception. They reference an overlapping, tiered approach to testing defined as:

- Tier 1: Cystic fibrosis, spinal muscular atrophy, and risk-based screening.
- Tier 2: “Conditions that have a severe or moderate phenotype and a carrier frequency of at least 1/100.” (Includes Tier 1).
- Tier 3: “Conditions with a carrier frequency $\geq 1/200$.” (Includes Tier 2). Includes X-linked conditions.
- Tier 4: “Genes less common than those in Tier 3 and can identify additional at-risk couples.” Conditions with a carrier frequency $<1/200$. (Includes Tier 3).

In terms of what screening approaches should be offered, the ACMG recommends that “all pregnant patients and those planning a pregnancy should be offered Tier 3 carrier screening. Tier 4 screening should be considered: when a pregnancy stems from a known or possible consanguineous relationship (second cousins or closer); [or] when a family or personal medical history warrants.” The ACMG does

NOT recommend “Offering Tier 1 and/or Tier 2 screening, because these do not provide equitable evaluation of all racial/ethnic groups [or] routine offering of Tier 4 panels.”

In terms of what autosomal recessive conditions are appropriate for carrier screening, the ACMG recommends that “All pregnant patients and those planning a pregnancy should be offered Tier 3 carrier screening for autosomal recessive and X-linked conditions. Reproductive partners of pregnant patients and those planning a pregnancy may be offered Tier 3 carrier screening for autosomal recessive conditions when carrier screening is performed simultaneously with their partner.”

In terms of which X-linked conditions are appropriate for carrier screening, they note that “All XX patients should be offered screening for only those X-linked genes [listed here] as part of Tier 3 screening.” The X-linked genes are: *ABCD1, AFF2, ARX, DMD, F8, F9, FMR1, GLA, L1CAM, MID1, NROB1, OTC, PLP1, PRGR, RS1, SLC6A8*.

Lastly, the ACMG notes the “critical” importance of education and counseling in carrier screening and recommends that “carrier screening counseling should be provided by knowledgeable and appropriately trained health-care professionals and should be performed pre- and post-test.”⁴²

British Fertility Society (BFS) Policy and Practice Guidelines

The BFS have published guidelines regarding PGS. These guidelines state that “It remains possible that PGS may be of benefit under certain circumstances. However at present patients should be informed that there is no robust evidence that PGS for advanced maternal age improves live birth rate per cycle started, and PGS should preferably be offered within the context of robustly designed randomised trials performed in suitably experienced centres.”⁴³

Indian Society for Assisted Reproduction (ISAR)

The Indian Society for Assisted Reproduction released consensus guidelines about preimplantation genetic testing in In-vitro-fertilization clinics in India. The recommendations for PGT-A are: “PGT-A is recommended for: advanced maternal age (36–40 years), or repeated pregnancy loss – known etiologies. PGT-A is not recommended for: young, good prognosis patients (<35 years), or unexplained RPL, or low AMH – limited eggs; multiple IVF cycles may be necessary in order to obtain one euploid blastocyst.”

The clinical recommendations for genetic testing for monogenic indications are:

- “Can be offered to all patients with single or multiple gene disorders with a positive mutation report
- Cannot be offered for diseases which as multifactorial and nongenetic based diseases
- In case PGT-A is desired, it should be performed on PGT-M screened embryos.”⁴⁴

VII. Applicable State and Federal Regulations

DISCLAIMER: If there is a conflict between this policy and any relevant, applicable government policy for a particular member [e.g., Local Coverage Determinations (LCDs) or National Coverage Determinations (NCDs) for Medicare and/or state coverage for Medicaid], then the government policy will be used to make the determination. For the most up-to-date Medicare policies and coverage, please visit the

Medicare search website: <https://www.cms.gov/medicare-coverage-database/search.aspx>. For the most up-to-date Medicaid policies and coverage, visit the applicable state Medicaid website.

Food and Drug Administration (FDA)

Many labs have developed specific tests that they must validate and perform in house. These laboratory-developed tests (LDTs) are regulated by the Centers for Medicare and Medicaid (CMS) as high-complexity tests under the Clinical Laboratory Improvement Amendments of 1988 (CLIA '88). LDTs are not approved or cleared by the U. S. Food and Drug Administration; however, FDA clearance or approval is not currently required for clinical use.

VIII. Applicable CPT/HCPCS Procedure Codes

CPT	Code Description
81228	Cytogenomic (genome-wide) analysis for constitution chromosomal abnormalities; interrogation of genomic regions for copy number variants, comparative genomic hybridization [CGH] microarray analysis
81229	Cytogenomic (genome-wide) analysis for consitutional chromosomal abnormalities; interrogation og genomic regions for copy number and single nucleotide polymorphism (SNP) variants, comparative genomic hybridization (CGH) microarray analysis
81349	Cytogenomic (genome-wide) analysis for constitutional chromosomal abnormalities; interrogation of genomic regions for copy number and loss-of-heterozygosity variants, low-pass sequencing analysis
81479	Unlisted molecular pathology procedure
0254U	Reproductive medicine (preimplantation genetic assessment), analysis of 24 chromosomes using embryonic DNA genomic sequence analysis for aneuploidy, and a mitochondrial DNA score in euploid embryos, results reported as normal (euploidy), monosomy, trisomy, or partial deletion/duplications, mosaicism, and segmental aneuploidy, per embryo tested Proprietary test: SMART PGT-A (Pre-implantation Genetic Testing - Aneuploidy) Lab/Manufacturer: Igenomix®

Current Procedural Terminology® American Medical Association. All Rights reserved.

Procedure codes appearing in Medical Policy documents are included only as a general reference tool for each policy. They may not be all-inclusive.

IX. Evidence-based Scientific References

- Schattman GL, Xu K. Preimplantation genetic testing. Updated January 19, 2024. <https://www.uptodate.com/contents/preimplantation-genetic-testing>
- PGDIS. Guidelines for good practice in PGD: programme requirements and laboratory quality assurance. *Reproductive biomedicine online*. Jan 2008;16(1):134-47.
- Stern HJ. Preimplantation Genetic Diagnosis: Prenatal Testing for Embryos Finally Achieving Its Potential. *J Clin Med*. 2014;3(1):280-309. doi:10.3390/jcm3010280
- Scott KL, Hong KH, Scott RT, Jr. Selecting the optimal time to perform biopsy for preimplantation genetic testing. *Fertility and sterility*. Sep 2013;100(3):608-14. doi:10.1016/j.fertnstert.2013.07.004
- Scott, Upham, Forman, Zhao, Treff NR. Cleavage-stage biopsy significantly impairs human embryonic implantation potential while blastocyst biopsy does not: a randomized and paired clinical trial. *Fertility and sterility*. Sep 2013;100(3):624-30. doi:10.1016/j.fertnstert.2013.04.039

6. Brezina PR, Anchan R, Kearns WG. Preimplantation genetic testing for aneuploidy: what technology should you use and what are the differences? *Journal of assisted reproduction and genetics*. Jul 2016;33(7):823-32. doi:10.1007/s10815-016-0740-2
7. ASRM. Use of preimplantation genetic diagnosis for serious adult onset conditions: a committee opinion. *Fertility and sterility*. Jul 2013;100(1):54-7. doi:10.1016/j.fertnstert.2013.02.043
8. ASRM. Use of reproductive technology for sex selection for nonmedical reasons. *Fertility and sterility*. Jun 2015;103(6):1418-22. doi:10.1016/j.fertnstert.2015.03.035
9. Vaiarelli A, Cimadomo D, Capalbo A, et al. Pre-implantation genetic testing in ART: who will benefit and what is the evidence? *Journal of assisted reproduction and genetics*. Oct 2016;33(10):1273-1278. doi:10.1007/s10815-016-0785-2
10. Alteri A, Corti L, Sanchez AM, Rabbellotti E, Papaleo E, Viganò P. Assessment of pre-implantation genetic testing for embryo aneuploidies: A SWOT analysis. *Clin Genet*. Apr 2019;95(4):479-487. doi:10.1111/cge.13510
11. Ledger W. Preimplantation genetic screening should be used in all in vitro fertilisation cycles in women over the age of 35 years: AGAINST: Pre-implantation genetic screening should not be used in all IVF cycles in women over the age of 35 years. *Bjog*. Dec 2019;126(13):1555. doi:10.1111/1471-0528.15942
12. Sullivan-Pyke C, Dokras A. Preimplantation Genetic Screening and Preimplantation Genetic Diagnosis. *Obstetrics and gynecology clinics of North America*. Mar 2018;45(1):113-125. doi:10.1016/j.ogc.2017.10.009
13. García-Herrero S, Martínez-Fernández A, Marin L, et al. New high-throughput semiautomated Next Generation Sequencing (NGS) platform for Pre-implantation Genetic Testing for aneuploidies (PGT-A). *Reproductive biomedicine online*. 2019;38(1)
14. Feldman B, Aizer A, Brengauz M, et al. Pre-implantation genetic diagnosis-should we use ICSI for all? *Journal of assisted reproduction and genetics*. Sep 2017;34(9):1179-1183. doi:10.1007/s10815-017-0966-7
15. Montazeri F, Foroughmand AM, Kalantar SM, Aflatoonian A, Khalilli MA. Tips and Tricks in Fluorescence In-situ Hybridization (FISH)-based Preimplantation Genetic Diagnosis/Screening (PGD/PGS). *International Journal of Medical Laboratory*. 2018;5(5):84-98.
16. Farra C, Choucair F, Awwad J. Non-invasive pre-implantation genetic testing of human embryos: an emerging concept. *Hum Reprod*. Dec 1 2018;33(12):2162-2167. doi:10.1093/humrep/dey314
17. Natera. Spectrum®. <https://www.natera.com/womens-health/spectrum-preimplantation-genetics/#pg-menu-tabs>
18. Simon AL, Kiehl M, Fischer E, et al. Pregnancy outcomes from more than 1,800 in vitro fertilization cycles with the use of 24-chromosome single-nucleotide polymorphism-based preimplantation genetic testing for aneuploidy. *Fertility and sterility*. Jul 1 2018;110(1):113-121. doi:10.1016/j.fertnstert.2018.03.026
19. iGLS. Preimplantation Genetic Testing for Aneuploidy. 2022;
20. PacGenomics. Preimplantation Genetic Testing (PGT). 2022;
21. RGI. What is PGT. 2022;
22. Dreesen J, Destouni A, Kourlaba G, et al. Evaluation of PCR-based preimplantation genetic diagnosis applied to monogenic diseases: a collaborative ESHRE PGD consortium study. *European journal of human genetics : EJHG*. Aug 2014;22(8):1012-8. doi:10.1038/ejhg.2013.277
23. Ghioffi CE, Goldberg JD, Haque IS, Lazarin GA, Wong KK. Clinical Utility of Expanded Carrier Screening: Reproductive Behaviors of At-Risk Couples. *Journal of genetic counseling*. Jun 2018;27(3):616-625. doi:10.1007/s10897-017-0160-1

24. Kamath MS, Antonisamy B, Sunkara SK. Zygotic splitting following embryo biopsy: a cohort study of 207 697 single-embryo transfers following IVF treatment. *Bjog*. Dec 12 2019;doi:10.1111/1471-0528.16045
25. Verpoest W, Staessen C, Bossuyt PM, et al. Preimplantation genetic testing for aneuploidy by microarray analysis of polar bodies in advanced maternal age: a randomized clinical trial. *Hum Reprod*. Sep 1 2018;33(9):1767-1776. doi:10.1093/humrep/dey262
26. Cornelisse S, Zagers M, Kostova E, Fleischer K, Wely M, Mastenbroek S. Preimplantation genetic testing for aneuploidies (abnormal number of chromosomes) in in vitro fertilisation. *Cochrane Database of Systematic Reviews*. 2020;(9)doi:10.1002/14651858.CD005291.pub3
27. Yap W, Lee C, Chan W, Lim Y. Detection of Mosaicism in Blastocyst using High Resolution Next Generation Sequencing Preimplantation Genetic Screening (hr-NGS). *Reproductive biomedicine online*. 2019;38(1)
28. Zeevi DA, Backenroth D, Hakam-Spector E, et al. Expanded clinical validation of Haploseek for comprehensive preimplantation genetic testing. *Genetics in Medicine*. 2021/07/01 2021;23(7):1334-1340. doi:10.1038/s41436-021-01145-6
29. Kumar A, Im K, Banjevic M, et al. Whole-genome risk prediction of common diseases in human preimplantation embryos. *Nat Med*. Mar 2022;28(3):513-516. doi:10.1038/s41591-022-01735-0
30. ACOG. Carrier Screening in the Age of Genomic Medicine. 2017;
31. ACOG. Preimplantation Genetic Testing. *Obstetrics & Gynecology*. 2020;135(3)
32. ACOG. ACOG Committee Opinion No. 410: Ethical issues in genetic testing. *Obstet Gynecol*. Jun 2008;111(6):1495-502. doi:10.1097/AOG.0b013e31817d252f
33. ASRM. Use of preimplantation genetic testing for monogenic defects (PGT-M) for adult-onset conditions: an Ethics Committee opinion. 2018;doi:10.1016/j.fertnstert.2018.04.003
34. Harton GL, De Rycke M, Fiorentino F, et al. ESHRE PGD consortium best practice guidelines for amplification-based PGD†. *Human Reproduction*. 2010;26(1):33-40. doi:10.1093/humrep/deq231
35. ESHRE PGT Consortium Steering Committee, Carvalho F, Coonen E, et al. ESHRE PGT Consortium good practice recommendations for the organisation of PGT†. *Human Reproduction Open*. 2020;2020(3)doi:10.1093/hropen/hoaa021
36. ASRM. Indications and management of preimplantation genetic testing for monogenic conditions: a committee opinion. *Fertility and sterility*. Jul 2023;120(1):61-71. doi:10.1016/j.fertnstert.2023.03.003
37. Ethics Committee of the American Society for Reproductive Medicine ASRM. Use of preimplantation genetic diagnosis for serious adult onset conditions: a committee opinion. *Fertility and sterility*. Jul 2013;100(1):54-7. doi:10.1016/j.fertnstert.2013.02.043
38. ASRM. Preimplantation genetic testing: a Practice Committee opinion. *Fertility and sterility*. Nov 2008;90(5 Suppl):S136-43. doi:10.1016/j.fertnstert.2008.08.062
39. ASRM. Use of preimplantation genetic testing for monogenic adult-onset conditions: an Ethics Committee opinion. <https://www.asrm.org/practice-guidance/ethics-opinions/use-of-preimplantation-genetic-testing-for-monogenic-adult-onset-conditions-an-ethics-committee-opinion/>
40. Leigh D, Cram DS, Rechitsky S, et al. PGDIS position statement on the transfer of mosaic embryos 2021. *Reproductive biomedicine online*. Jul 2022;45(1):19-25. doi:10.1016/j.rbmo.2022.03.013
41. Grody WW, Thompson BH, Gregg AR, et al. ACMG position statement on prenatal/preconception expanded carrier screening. *Genet Med*. Jun 2013;15(6):482-3. doi:10.1038/gim.2013.47
42. Gregg AR, Aarabi M, Klugman S, et al. Screening for autosomal recessive and X-linked conditions during pregnancy and preconception: a practice resource of the American College of Medical

Genetics and Genomics (ACMG). *Genet Med.* Oct 2021;23(10):1793-1806. doi:10.1038/s41436-021-01203-z

43. Anderson RA, Pickering S. The current status of preimplantation genetic screening: British Fertility Society Policy and Practice Guidelines. *Hum Fertil (Camb)*. Jun 2008;11(2):71-5. doi:10.1080/14647270802041607
44. Malhotra J, Malhotra K, Majumdar G, et al. Indian Society for Assisted Reproduction Consensus Guidelines on Preimplantation Genetic Testing in In vitro Fertilization Clinics. *J Hum Reprod Sci.* Nov 2021;14(Suppl 1):S31-s47. doi:10.4103/0974-1208.330503

X. Review/Revision History

Effective Date	Summary
07/01/2025	Reviewed and Updated: Updated background, guidelines, and evidence-based scientific references. Literature review necessitated the following changes in coverage criteria: New CC2.g. to allow PGT-M for Huntington and ADPKD: “g) One biological parent is a known carrier of one of the following adult-onset, autosomal dominant disorders: i) Huntington Disease (HTT) ii) Autosomal dominant kidney disease (PKD, PKD2)” With the change to CC2, added “not discussed above” to CC4.a. Now reads: “a) Preimplantation genetic testing for adult-onset disorders not discussed above.”
12/01/2024	Initial Policy Implementation