

Genetic Testing for Lactase Insufficiency

Policy Number: AHS – M2080 – Genetic Testing for Lactase Insufficiency	Policy Revision Date: 10/15/2025 Initial Policy Effective Date: 12/01/2024
--	---

[POLICY DESCRIPTION](#) | [RELATED POLICIES](#) | [INDICATIONS AND/OR LIMITATIONS OF COVERAGE](#) | [TABLE OF TERMINOLOGY](#) | [SCIENTIFIC BACKGROUND](#) | [GUIDELINES AND RECOMMENDATIONS](#) | [APPLICABLE STATE AND FEDERAL REGULATIONS](#) | [APPLICABLE CPT/HCPCS PROCEDURE CODES](#) | [EVIDENCE-BASED SCIENTIFIC REFERENCES](#) | [REVISION HISTORY](#)

I. Policy Description

Lactose intolerance (LI) is a common clinical syndrome defined by abdominal pain, flatulence, bloating, borborygmus and osmotic diarrhea; LI is caused by the breakdown of nondigested lactose by the gut microflora.¹

Lactose malabsorption (LM) is the non-digestion of lactose caused by low expression of the enzyme lactase and is a physiological feature occurring in most mammals after infancy.^{1,2}

II. Related Policies

Policy Number	Policy Title
N/A	Not Applicable

III. Indications and/or Limitations of Coverage

Application of coverage criteria is dependent upon an individual’s benefit coverage at the time of the request. Specifications pertaining to Medicare and Medicaid can be found in the “Applicable State and Federal Regulations” section of this policy document.

The following does not meet coverage criteria due to a lack of available published scientific literature confirming that the test(s) is/are required and beneficial for the diagnosis and treatment of an individual’s illness.

- 1) For the prediction of lactase insufficiency, targeted mutation analysis of *LCT* -13910 C>T **DOES NOT MEET COVERAGE CRITERIA.**
- 2) For individuals with lactose intolerance and/or lactase insufficiency, genetic testing of *LCT* and/or *MCM6* **DOES NOT MEET COVERAGE CRITERIA.**

IV. Table of Terminology

Term	Definition
AAP	American Academy of Pediatrics
ACG	American College of Gastroenterology

BSACI	British Society for Allergy & Clinical Immunology
C/T-13910	Eurasian lactase persistence variant
CLIA	Clinical Laboratory Improvement Amendments
CMS	Centers for Medicare and Medicaid Services
FODMAP	Fermentable oligo-, di- and monosaccharides and polyols
FGID	Functional gastrointestinal disorders
HBT	Hydrogen breath test
IBD	Inflammatory bowel disease
LBT	Lactose breath test
LCT	Lactase/lactase-phlorizin hydrolase
LD	Lactase deficiency
LDTs	Laboratory developed tests
LI	Lactose intolerance
LIP	Lactose-intolerant phenotype
LM	Lactose malabsorption
LNP	Lactase non-persistence
LTT	Lactose tolerance test
MCM6	<i>Minichromosome maintenance complex component 6</i>
NIH	National Institutes of Health
PCR	Polymerase chain reaction
RNA	Ribonucleic acid
RT-PCR	Real-time polymerase chain reaction
SNPs	Single nucleotide polymorphisms

V. Scientific Background

Lactose is a disaccharide, which is a class of sugars comprised of two monosaccharides or simple sugars. Lactose consists of galactose bound to glucose and is the main source of carbohydrates from mammalian milk. Intestinal absorption of lactose requires initial hydrolysis by the enzyme lactase. Low lactase activity results in undigested lactose and colonic bacterial fermentation of that lactose. This leads to the characteristic symptoms of LI, such as bloating and flatulence.³

Lactase expression decreases as consequence of the normal maturational down-regulation after weaning, ultimately to undetectable levels in most populations.⁴ Lactase expression persists, however, in descendants of populations that traditionally practice cattle domestication who maintain the ability to digest milk and other dairy products into adulthood.⁵ Adult expression of the gene encoding lactase (LCT), located on chromosome two at 2q21 appears to be regulated by *cis*-acting elements⁶ and is inherited as an autosomal recessive trait.⁷ The LCT gene is regulated by the nearby *MCM6* gene (minichromosome maintenance complex component 6), which encodes a helicase complex. A few different *MCM6* variants have been identified in affecting LCT gene expression.⁸ Single nucleotide polymorphisms (SNPs) associated with the lactase persistence vary by region. In European populations it is associated with C/T-13910 and G/A-22018 mutations of the LCT gene.^{7,9-11} Additional mutations have been identified in Saudi Arabia populations with G-13915,¹² and in African tribes with the G-14010, G-13915, and G-13907 polymorphisms.^{13,14} No SNP associated with lactase persistence has been identified


in the lactase gene regulatory sequence in Chinese populations.¹⁵ In adult patients with homozygous lactase persistence, enzyme levels are 10-times higher than for patients with homozygous non-persistence, and heterozygous individuals.^{5,16}

Lactase deficiency Baffour-Awuah, et al. (2015) is defined as markedly reduced brush-border lactase activity relative to the activity observed in infants.⁵ Continued dairy consumption despite low expression of lactase results in unabsorbed lactose being present in the intestinal tract (lactose malabsorption [LM]), which can lead to symptoms of lactose intolerance (LI) in susceptible individuals.¹⁸

Lactose intolerance is defined by patient reports of abdominal pain, bloating, borborygmi, and diarrhea induced by dairy consumption. Unabsorbed lactose increases the osmotic load thus increasing the intestinal water content, resulting in osmotic diarrhea. Additionally, lactose and other poorly-absorbed oligosaccharides, disaccharides, monosaccharides, and polyols ubiquitous in the diet are readily fermented by the colonic microbiome, leading to production of short-chain fatty acids and gas (mainly hydrogen [H₂], carbon dioxide [CO₂], and methane [CH₄]).^{19,20} LI may be associated with nonspecific symptoms, abdominal pain, bloating, flatulence, diarrhea, or vomiting.¹⁸ however, it is unclear whether these symptoms are directly due to lactose ingestion. Although LM is nearly always attributable to LD, it is not possible to make a definitive diagnosis on clinical presentation alone because double-blind trials have shown that the reliability of self-reported LI is very poor.^{5,15,21}

Determining if reported symptoms of LI are resultant from LD can be approached through several different methods. The gold standard is the measurement of lactase, sucrase, and maltase activity through intestinal biopsies. However, this method is not commonly used due to its invasive nature.² Other tests, such as the lactose breath test or biochemical blood tests are more frequently used.^{1,22,23} In addition to biochemical blood tests, genetic markers may be useful for LI diagnosis; however, a positive genetic test indicates whether lactase activity decline may represent a clinical problem for the patient, but the test does not give information on actual patient symptoms, making it inappropriate as an initial screening as not all patients with LM will develop symptoms of LI.¹

On the contrary, this information is more readily accessible by combining the lactose breath test with intolerance symptom evaluation.²⁴ Usual LI management involves excluding milk and milk products from the diet, while ensuring adequate calcium intake.^{1,25,26} The use of genetic tests has been proposed as an adjunct to LI diagnosis to differentiate primary LD from secondary causes²⁷ as depicted below in the figure taken from Ponte, et al. (2016).



Clinical Utility and Validity

Marton, et al. (2012) compared the common polymorphism C/T 13910 with the lactase breath test and lactose tolerance test to assess each test’s ability to predict genotype/phenotype relationships. The agreement of the breath test and genotype was 0.88 sensitivity and 0.85 specificity whereas the agreement between genotype and tolerance test was 0.94 sensitivity and 0.90 specificity.²⁸

Baffour-Awuah, et al. (2015) studied the association of genotypes at the -13910 and -22018 SNPs with clinical characteristics, RNA quantification, and enzymatic phenotypes among a range of European ethnicities within the U.S. population. The authors concluded that “13910T/T genotype will frequently, but not perfectly, predict lactase persistence in this mixed European-ancestry population; a -13910T/C genotype will not predict the phenotype.”¹⁷

Misselwitz, et al. (2013) stated that genetic testing for the -13910*T genotype in certain African, Arabic, or Asian subpopulations has limited value because lactase persistence may be linked to different polymorphisms. The authors also stated that genetic tests will be negative in patients with secondary causes of lactase deficiency and that no information about clinical symptoms LI will be obtained during testing.²⁹

Brasen, et al. (2017) genotyped 3395 routine samples using real-time polymerase chain reaction (PCR) for the -13910C > T variant to determine the prevalence of the variants in a Danish cohort examined for lactose intolerance as well as to improve the real-time PCR analysis for detection of the different variants. The researchers found that "Using real-time PCR resulted in 100% successful genotyping of the -13910C > T variant. By using a quality value of 99% and sequencing the undetermined samples we improved the ability of the assay to identify variants other than -13910C > T. This resulted in a reduction of the diagnostic error rate by a factor of 2.4 while increasing the expenses only 3%."³⁰

Coluccia, et al. (2019) compared results from a lactose breath test and a lactase gene polymorphism test in a total of 158 symptomatic adults to identify lactose intolerance. Whole blood samples were used for genetic testing purposes. Lactose breath testing resulted in positive results in 75.9% of participants, while genetic testing identified lactase gene polymorphisms in 82.3% of participants. In conclusion, the authors state, "We suggest considering the use of the genetic test after LBT [lactose breath test] administration, when secondary hypolactasia is suspected, for completion of diagnostic procedures."³¹

Muendlein, et al. (2019) developed a real-time PCR protocol for the detection of the LCT-13910C>T variant from whole blood samples. Lactose tolerance in adults is known to be strongly associated with the genetic variant LCT-13910C>T, which lies near the lactase encoding gene. Results from real-time PCR were compared to those from Sanger sequencing. A total of 105 whole blood samples were analyzed. When compared to Sanger sequencing, the real-time PCR protocol was determined to be a reliable method for the detection of the LCT-13910C>T.³²

In a cross-sectional study by Couce, et al. (2020), 493 children suspected of functional gastrointestinal disorders (FGID) were studied using the exhaled H₂ test, gastrointestinal symptoms, and SNP C/T-13910 genetic testing to observe the correlation between the genotype and phenotype of the C/T-13910 lactase non-persistence (LNP) gene. "The C/T-13910 genotype distribution was as follows: CC, 46.0%; CT, 39.5%; TT 14.4%. The frequency of the LP allele was 34.1%." A significant increase in LNP genotype, H₂ test, and gastrointestinal symptoms were observed with increasing age. According to phenotype, CC carriers were mainly lactose non-absorbers (75.4%), TT carriers were lactose absorbers (91.6%), and CT heterozygotes were predominantly lactose absorbers (90.7%) and lactose tolerant (74.3%). According to these results and its prevalence in older age, the authors state that the C/T-13910 polymorphism is significantly correlated with the phenotype and the authors suggest that the practical value of genetic testing is greater in older children.³³

Nardone, et al. (2021) studied the prevalence of LI assessed by hydrogen breath test (H-BT) and investigated the prevalence of three single genetic polymorphisms of the lactase gene. A total of 54 IBD patients were recruited, and H-BT was positive in 64.8% of IBD patients and 62.3% control patients. The genetic analysis revealed that 46 IBD patients (85.2%) had the wild-type genotype (LCT-13910 CC) while the other polymorphisms (CT-22018, AG-13910, and CT-22018/AG-13910) were less common in IBD patients. In the control group, the wild genotype was found in 87% of the patients. Therefore, the prevalence of polymorphisms did not differ between the IBD group and the control group. The correlation between positive H-BT and genetic analysis showed that the "wild-type genotype was associated with higher rate of lactose intolerance in the total population."³⁴

Usai-Satta, et al. (2021) reviewed the usefulness of hydrogen breath testing in the nutritional management of carbohydrate digestive disorders, including LM and intolerance, using the online databases PubMed, Medline, and Cochrane. According to the authors, the literature search yielded that though lactose breath testing is an indirect test for LM, "lactose BT showed good sensitivity and optimal

specificity for lactose malabsorption”, leading them to conclude that “before starting a low FODMAP [fermentable oligo-, di- and monosaccharides and polyols] diet, lactose BT should be suggested in a population with low prevalence of hypolactasia,” helped in part by its non-invasiveness and inexpensiveness.³⁵

Cavichio, et al. (2024) studied the agreement between C/T-13910 polymorphism genotyping results and lactose tolerance test (LTT) results in Brazil. LTT is the most commonly used LI diagnostic test in Brazil. The C/T-13910 polymorphism has mostly been studied in Caucasian populations; therefore, the authors of this study aimed to investigate the accuracy of C/T-13910 polymorphism genotyping in Brazil’s “highly mixed” population. The authors included 404 patients who had undergone both C/T-13910 polymorphism genotyping and an LTT. Overall, “there was agreement between the genotyping and LTT results in 325 (80.4%) patients and discordance in 79 (19.6%) patients.” The authors classified these results as “moderate agreement” and recommend additional studies to further investigate.³⁶

VI. Guidelines and Recommendations

American Academy of Pediatrics (AAP)

The AAP published guidelines³⁷ on the evaluation of Lactose Intolerance in Infants, Children and Adolescents which recommend:

“Children with suspected lactose intolerance can be assessed clinically by dietary lactose elimination or by tests including noninvasive hydrogen breath testing or invasive intestinal biopsy determination of lactase (and other disaccharidase) concentrations. Treatment consists of use of lactase-treated dairy products or oral lactase supplementation, limitation of lactose-containing foods, or dairy elimination... If dairy products are eliminated, other dietary sources of calcium or calcium supplements need to be provided.”

The AAP also reported that “Recent studies suggest that in the future, genetic testing may be useful for identifying individuals at increased risk of lactase deficiency and consequent diminished bone mineral density, potentially allowing early intervention with dietary manipulation or nutrient supplementation.”³⁷

This statement was reaffirmed in 2012.

American College of Gastroenterology (ACG)

The ACG has a webpage focused on LI in children. Regarding diagnoses, the ACG states that “lactose intolerance is diagnosed by a simple test called a hydrogen breath test.”³⁸ Genetic testing was not mentioned.

The 1st Rome Consensus Conference

The 1st Rome Consensus Conference issued their opinion on H₂-breath testing for sugar malabsorption. They found that “the determination of lactase activity in jejunal biopsy is currently considered the gold standard for lactose malabsorption. However, its results can be influenced by the irregular dissemination of lactase activity throughout the small intestine mucosa. Based on literature review, the lactose breath test is a reliable, non-invasive technique, which is provided with good sensitivity and optimal specificity.”

Though it is acknowledged that there is no unequivocal reference test available for LM testing, the Conference graded the quality of evidence for the above as Class I (“Conditions with evidence or general accord that a particular procedure or treatment is useful or effective”) of Strength A (“Data derived from multiple large and intermediate RCT”).³⁹

VII. Applicable State and Federal Regulations

DISCLAIMER: If there is a conflict between this policy and any relevant, applicable government policy for a particular member [e.g., Local Coverage Determinations (LCDs) or National Coverage Determinations (NCDs) for Medicare and/or state coverage for Medicaid], then the government policy will be used to make the determination. For the most up-to-date Medicare policies and coverage, please visit the Medicare search website: <https://www.cms.gov/medicare-coverage-database/search.aspx>. For the most up-to-date Medicaid policies and coverage, visit the applicable state Medicaid website.

Food and Drug Administration (FDA)

Many labs have developed specific tests that they must validate and perform in house. These laboratory developed tests (LDTs) are regulated by the Centers for Medicare and Medicaid (CMS) as high-complexity tests under the Clinical Laboratory Improvement Amendments of 1988 (CLIA '88). LDTs are not approved or cleared by the U. S. Food and Drug Administration; however, FDA clearance or approval is not currently required for clinical use.

VIII. Applicable CPT/HCPCS Procedure Codes

CPT	Code Description
81400	Molecular pathology procedure, Level 1 (eg, identification of single germline variant [eg, SNP] by techniques such as restriction enzyme digestion or melt curve analysis)
81479	Unlisted molecular pathology procedure

Current Procedural Terminology[®] American Medical Association. All Rights reserved.

Procedure codes appearing in Medical Policy documents are included only as a general reference tool for each policy. They may not be all-inclusive.

IX. Evidence-based Scientific References

1. Ponte PR, de Medeiros PH, Havt A, et al. Clinical evaluation, biochemistry and genetic polymorphism analysis for the diagnosis of lactose intolerance in a population from northeastern Brazil. *Clinics (Sao Paulo, Brazil)*. Feb 2016;71(2):82-9. doi:10.6061/clinics/2016(02)06
2. Di Rienzo T, D'Angelo G, D'Aversa F, et al. Lactose intolerance: from diagnosis to correct management. *European review for medical and pharmacological sciences*. 2013;17 Suppl 2:18-25. <https://www.europeanreview.org/wp/wp-content/uploads/018-025.pdf>
3. Luyt D, Ball H, Makwana N, et al. BSACI guideline for the diagnosis and management of cow's milk allergy. *Clinical & Experimental Allergy*. May 01, 2014 2014;44(5):642-672. doi:10.1111/cea.12302
4. Swallow DM. Genetics of lactase persistence and lactose intolerance. *Annual review of genetics*. 2003;37:197-219. doi:10.1146/annurev.genet.37.110801.143820
5. Deng Y, Misselwitz B, Dai N, Fox M. Lactose Intolerance in Adults: Biological Mechanism and Dietary Management. *Nutrients*. 2015;7(9):8020-35. doi:10.3390/nu7095380

6. Wang Y, Harvey CB, Pratt WS, et al. The lactase persistence/non-persistence polymorphism is controlled by a cis-acting element. *Human molecular genetics*. Apr 1995;4(4):657-62. doi:10.1093/hmg/4.4.657
7. Enattah NS, Sahi T, Savilahti E, Terwilliger JD, Peltonen L, Jarvela I. Identification of a variant associated with adult-type hypolactasia. *Nature genetics*. Feb 2002;30(2):233-7. doi:10.1038/ng826
8. NIH. MCM6 gene. NIH. Updated March 24, 2023. <https://ghr.nlm.nih.gov/gene/MCM6>
9. Poulter M, Hollox E, Harvey CB, et al. The causal element for the lactase persistence/non-persistence polymorphism is located in a 1 Mb region of linkage disequilibrium in Europeans. *Annals of human genetics*. Jul 2003;67(Pt 4):298-311. doi:10.1046/j.1469-1809.2003.00048.x
10. Hogenauer C, Hammer HF, Mellitzer K, Renner W, Krejs GJ, Toplak H. Evaluation of a new DNA test compared with the lactose hydrogen breath test for the diagnosis of lactase non-persistence. *European journal of gastroenterology & hepatology*. Mar 2005;17(3):371-6. doi:10.1097/00042737-200503000-00018
11. Ridfelt P, Hakansson LD. Lactose intolerance: lactose tolerance test versus genotyping. *Scandinavian journal of gastroenterology*. Jul 2005;40(7):822-6. doi:10.1080/00365520510015764
12. Imtiaz F, Savilahti E, Sarnesto A, et al. The T/G 13915 variant upstream of the lactase gene (LCT) is the founder allele of lactase persistence in an urban Saudi population. *Journal of medical genetics*. Oct 2007;44(10):e89. doi:10.1136/jmg.2007.051631
13. Ingram CJ, Elamin MF, Mulcare CA, et al. A novel polymorphism associated with lactose tolerance in Africa: multiple causes for lactase persistence? *Human genetics*. Feb 2007;120(6):779-88. doi:10.1007/s00439-006-0291-1
14. Tishkoff SA, Reed FA, Ranciaro A, et al. Convergent adaptation of human lactase persistence in Africa and Europe. *Nature genetics*. 2007;39(1):31-40. doi:10.1038/ng1946
15. Zheng X, China ZUDoGH, Chu H, et al. Self-reported lactose intolerance in clinic patients with functional gastrointestinal symptoms: prevalence, risk factors, and impact on food choices. *Neurogastroenterology & Motility*. 2016;27(8):1138-1146. doi:10.1111/nmo.12602
16. Enattah NS, Kuokkanen M, Forsblom C, et al. Correlation of intestinal disaccharidase activities with the C/T-13910 variant and age. *World journal of gastroenterology*. 2007;13(25):3508-12. doi:10.3748/wjg.v13.i25.3508
17. Baffour-Awuah NY, Fleet S, Montgomery RK, et al. Functional significance of single nucleotide polymorphisms in the lactase gene in diverse US patients and evidence for a novel lactase persistence allele at -13909 in those of European ancestry. *Journal of pediatric gastroenterology and nutrition*. 2015;60(2):182-91. doi:10.1097/mpg.0000000000000595
18. Hammer HF, Högenauer C. Lactose intolerance and malabsorption: Clinical manifestations, diagnosis, and management. Updated October 01, 2024. <https://www.uptodate.com/contents/lactose-intolerance-and-malabsorption-clinical-manifestations-diagnosis-and-management>
19. Magge S, Lembo A. Low-FODMAP Diet for Treatment of Irritable Bowel Syndrome. *Gastroenterology & hepatology*. 2012;8(11):739-45.
20. Shepherd SJ, Lomer MC, Gibson PR. Short-chain carbohydrates and functional gastrointestinal disorders. *The American journal of gastroenterology*. May 2013;108(5):707-17. doi:10.1038/ajg.2013.96
21. Suarez FL, Savaiano DA, Levitt MD. A comparison of symptoms after the consumption of milk or lactose-hydrolyzed milk by people with self-reported severe lactose intolerance. *The New England journal of medicine*. 1995;333(1):1-4. doi:10.1056/nejm199507063330101
22. Furnari M, Bonfanti D, Parodi A, et al. A comparison between lactose breath test and quick test on duodenal biopsies for diagnosing lactase deficiency in patients with self-reported lactose

- intolerance. *Journal of clinical gastroenterology*. Feb 2013;47(2):148-52.
doi:10.1097/MCG.0b013e31824e9132
23. Mattar R, Basile-Filho A, Kemp R, Santos JS. Comparison of Quick Lactose Intolerance Test in duodenal biopsies of dyspeptic patients with single nucleotide polymorphism LCT-13910C>T associated with primary hypolactasia/lactase-persistence. *Acta cirurgica brasileira*. 2013;28 Suppl 1:77-82. doi:10.1590/s0102-86502013001300015
 24. Di Stefano M, Terulla V, Tana P, Mazzocchi S, Romero E, Corazza GR. Genetic test for lactase non-persistence and hydrogen breath test: is genotype better than phenotype to diagnose lactose malabsorption? *Digestive and liver disease : official journal of the Italian Society of Gastroenterology and the Italian Association for the Study of the Liver*. 2009;41(7):474-9.
doi:10.1016/j.dld.2008.09.020
 25. Misselwitz B. Lactose intolerance: new insights due to blinded testing? *Digestion*. 2014;90(1):72-3.
doi:10.1159/000365144
 26. Usai-Satta P, Scarpa M, Oppia F, Cabras F. Lactose malabsorption and intolerance: What should be the best clinical management? *World journal of gastrointestinal pharmacology and therapeutics*. Jun 6 2012;3(3):29-33. doi:10.4292/wjgpt.v3.i3.29
 27. Bodlaj G, Stocher M, Hufnagl P, et al. Genotyping of the lactase-phlorizin hydrolase -13910 polymorphism by LightCycler PCR and implications for the diagnosis of lactose intolerance. *Clinical chemistry*. 2006;52(1):148-51. doi:10.1373/clinchem.2005.057240
 28. Marton A, Xue X, Szilagyi A. Meta-analysis: the diagnostic accuracy of lactose breath hydrogen or lactose tolerance tests for predicting the North European lactase polymorphism C/T-13910. *Alimentary pharmacology & therapeutics*. Feb 2012;35(4):429-40. doi:10.1111/j.1365-2036.2011.04962.x
 29. Misselwitz B, Pohl D, Fruhauf H, Fried M, Vavricka SR, Fox M. Lactose malabsorption and intolerance: pathogenesis, diagnosis and treatment. *United European gastroenterology journal*. 2013;1(3):151-9. doi:10.1177/2050640613484463
 30. Brasen CL, Frischknecht L, Ornskov D, Andreassen L, Madsen JS. Combination of real-time PCR and sequencing to detect multiple clinically relevant genetic variations in the lactase gene. *Scandinavian journal of clinical and laboratory investigation*. 2017;77(1):60-65.
doi:10.1080/00365513.2016.1261408
 31. Coluccia E, Iardino P, Pappalardo D, et al. Congruency of Genetic Predisposition to Lactase Persistence and Lactose Breath Test. *Nutrients*. 2019;11(6)doi:10.3390/nu11061383
 32. Muendlein A, Leiherer A, Zach C, et al. Real-time PCR based detection of the lactase non-persistence associated genetic variant LCT-13910C>T directly from whole blood. *Mol Biol Rep*. 2019;46(2):2379-2385. doi:10.1007/s11033-019-04696-9
 33. Couce ML, Sánchez-Pintos P, González-Vioque E, Leis R. Clinical Utility of LCT Genotyping in Children with Suspected Functional Gastrointestinal Disorder. *Nutrients*. 2020;12(10):3017.
doi:10.3390/nu12103017
 34. Nardone OM, Manfellotto F, D'Onofrio C, et al. Lactose Intolerance Assessed by Analysis of Genetic Polymorphism, Breath Test and Symptoms in Patients with Inflammatory Bowel Disease. *Nutrients*. 2021;13(4):1290. doi:10.3390/nu13041290
 35. Usai-Satta P, Oppia F, Lai M, Cabras F. Hydrogen Breath Tests: Are They Really Useful in the Nutritional Management of Digestive Disease? *Nutrients*. Mar 17 2021;13(3)doi:10.3390/nu13030974
 36. Cavichio MWE, Quaió C, Baratela W, Oliveira PMC, Tahan S. Evaluation of Agreement Between C/T-13910 Polymorphism Genotyping Results and Lactose Tolerance Test Results: A Retrospective

Population-Based Study in Brazil. *Arq Gastroenterol.* 2024;61:e23104. doi:10.1590/s0004-2803.24612023-104

- 37. Heyman MB. Lactose intolerance in infants, children, and adolescents. *Pediatrics.* Sep 2006;118(3):1279-86. doi:10.1542/peds.2006-1721
- 38. ACG. Lactose Intolerance. <https://gi.org/topics/lactose-intolerance-in-children/>
- 39. Gasbarrini A, Corazza GR, Gasbarrini G, et al. Methodology and indications of H2-breath testing in gastrointestinal diseases: the Rome Consensus Conference. *Alimentary pharmacology & therapeutics.* Mar 30 2009;29 Suppl 1:1-49. doi:10.1111/j.1365-2036.2009.03951.x

X. Review/Revision History

Effective Date	Summary
10/15/2025	Reviewed and Updated: Updated the background, guidelines and recommendations, and evidence-based scientific references. Literature review did not necessitate any modifications to coverage criteria.
12/01/2024	Reviewed and Updated: Updated the background, guidelines and recommendations, and evidence-based scientific references. Literature review did not necessitate any modifications to coverage criteria.
12/01/2024	Initial Policy Implementation