

## Identification of Microorganisms using Nucleic Acid Probes

Policy Number: AHS – M2097 – Identification of Microorganisms Using Nucleic Acid Probes	Prior Policy Name and Number, as applicable:
Initial Presentation Date: 06/01/2021 Revision Date: N/A	

### I. Policy Description

Nucleic acid hybridization technologies utilize complementary properties of the DNA double-helix structures to anneal together DNA fragments from different sources. These techniques are utilized in polymerase chain reaction (PCR) and fluorescent resonance energy transfer (FRET) techniques to identify microorganisms (Khan, 2014).

### II. Related Policies

Policy Number	Policy Title
AHS-G2143	Lyme Disease
AHS-G2149	Pathogen Panel Testing
AHS-G2157	Diagnostic Testing of Common Sexually Transmitted Infections
AHS-G2158	Testing for Mosquito- or Tick-Related Infections
AHS-M2057	Diagnosis of Vaginitis Including Multi-Target PCR Testing

### III. Indications and/or Limitations of Coverage

*Application of coverage criteria is dependent upon an individual’s benefit coverage at the time of the request. Medical Policy Statements do not ensure an authorization or payment of services. Please refer to the plan contract (often referred to as the Evidence of Coverage) for the service(s) referenced in the Medical Policy Statement. If there is a conflict between the Medical Policy Statement and the plan contract (i.e., Evidence of Coverage), then the plan contract (i.e., Evidence of Coverage) will be the controlling document used to make the determination.*

*Application of coverage criteria is dependent upon an individual’s benefit coverage at the time of the request. If there is a conflict between this Policy and any relevant, applicable government policy [e.g. National Coverage Determinations (NCDs) for Medicare] for a particular member, then the government policy will be used to make the determination. For the most up-to-date Medicare policies and coverage,*

please visit their search website <http://www.cms.gov/medicare-coverage-database/overview-and-quick-search.aspx?from2=search1.asp&> or the manual website

1. The status of nucleic acid identification using direct probe, amplified probe, or quantification for the microorganism’s procedure codes is summarized in Table 1 below. "MCC" in the table below indicates that the test **MEETS COVERAGE CRITERIA**; while “DNMCC” tests indicates that the test **DOES NOT MEET COVERAGE CRITERIA**.

Microorganism	Direct Probe	Amplified Probe	Quantification
<i>Bartonella henselae</i> or <i>quintana</i>		87471(MCC)	87472 (DNMCC)
<i>Candida</i> species (For vaginitis, please review AHS-M2057 Diagnosis of Vaginitis)	87480 (MCC) for vaginitis	87481 (DNMCC) for all situations	87482 (DNMCC) for all situations
Including Multi-Target PCR Testing)	87480 (DNMCC) for all other situations except vaginitis		
<i>Chlamydia pneumoniae</i>	87485 (MCC)	87486 (MCC)	87487 (DNMCC)
<i>Clostridium difficile</i>	87493 (MCC)		
<i>Cytomegalovirus</i>	87495 (MCC)	87496 (MCC)	87497 (MCC)
<i>Enterococcus</i> , Vancomycin-resistant (e.g., enterococcus vanA, vanB)		87500 (MCC)	
<i>Enterovirus</i>		87498 (MCC)	
Hepatitis B		87516 (MCC)	87517 (MCC)
Hepatitis G	87525 (DNMCC)	87526 (DNMCC)	87527 (DNMCC)
Herpes virus-6	87531 (MCC)	87532 (DNMCC)	87533 (MCC)
<i>Legionella pneumophila</i>	87540 (MCC)	87541 (MCC)	87542 (DNMCC)
<i>Mycoplasma pneumoniae</i>	87580 (MCC)	87581 (MCC)	87582 (DNMCC)
<i>Mycoplasma genitalium</i>		87563 (MCC)	
Respiratory syncytial virus		87634 (MCC)	
<i>Staphylococcus aureus</i>		87640 (MCC)	

<i>Staphylococcus aureus</i> , methicillin resistant		87641 (MCC)	
---	--	-------------	--

\*DNMCC= Does Not Meet Coverage Criteria; MCC= Meets coverage criteria.

2. The technique for quantification includes both amplification and direct probes; therefore, simultaneous coding for both amplification or direct probes **DOES NOT MEET COVERAGE CRITERIA**.
3. PCR testing for the following microorganisms that do not have specific CPT codes **MEETS COVERAGE CRITERIA** (not an all-inclusive list):
  - a. Actinomyces, for identification of actinomyces species in tissue specimens
  - b. Adenovirus, to diagnose adenovirus myocarditis, and to diagnose adenovirus infection in immunocompromised hosts, including transplant recipients
  - c. *Bacillus Anthracis*
  - d. BK polyomavirus in transplant recipients receiving immunosuppressive therapies and persons with immunosuppressive diseases
  - e. *Bordetella pertussis* and *B. parapertussis*, for diagnosis of whooping cough in individuals with coughing
  - f. *Brucella spp.*, for members with signs and symptoms of Brucellosis, and history of direct contact with infected animals and their carcasses or secretions or by ingesting unpasteurized milk or milk products
  - g. *Burkholderia* infections (including *B. cepacia*, *B. gladioli*), diagnosis
  - h. Chancroid (*Haemophilus ducreyi*), for diagnosis of persons with genital ulcer disease
  - i. *Coxiella burnetii*, for confirmation of acute Q fever
  - j. EBOLA
  - k. Epidemic typhus (*Rickettsia prowazekii*), diagnosis
  - l. Epstein Barr Virus (EBV): for detection of EBV in post-transplant lymphoproliferative disorder; or for testing for EBV in persons with lymphoma; or for those who are immunocompromised for other reasons.
  - m. *Francisella tularensis*, for presumptive diagnosis of tularemia
  - n. Hantavirus, diagnosis
  - o. Hemorrhagic fevers and related syndromes caused by viruses of the family *Bunyaviridae*

(Rift Valley fever, Crimean-Congo hemorrhagic fever, hemorrhagic fever with renal syndromes), for diagnosis in acute phase in persons with clinical presentation suggestive of these conditions

- p. Hepatitis D virus, for confirmation of active infection in persons with anti-HDV antibodies
- q. Hepatitis E virus, for definitive diagnosis in persons with anti-HEV antibodies
- r. Human T Lymphotropic Virus type 1 and type 2 (HTLV-I and HTLV-II), to confirm the presence of HTLV-I and HTLV-II in the cerebrospinal fluid of persons with signs or symptoms of HTLV-I/HTLV-II
- s. Human metapneumovirus
- t. JC polyomavirus, in transplant recipients receiving immunosuppressive therapies, in persons with immunosuppressive diseases, and for diagnosing progressive multifocal leukoencephalopathy in persons with multiple sclerosis or Crohn's disease receiving natalizumab (Tysabri)
- u. Leishmaniasis, diagnosis
- v. Measles virus (Morbilliviruses), for diagnosis of measles
- w. Mumps
- x. *Neisseria meningitidis*, to establish diagnosis where antibiotics have been started before cultures have been obtained
- y. Parvovirus, for detecting chronic infection in immunocompromised persons
- z. Psittacosis, for diagnosis of *Chlamydomphila (Chlamydia) psittaci* infection
  - aa. Rubella, diagnosis
  - bb. *Toxoplasma gondii*, for detection of T. gondii infection in immunocompromised persons with signs and symptoms of toxoplasmosis, and for detection of congenital *Toxoplasma gondii* infection (including testing of amniotic fluid for toxoplasma infection)
  - cc. Varicella-Zoster infections
  - dd. Whipple's disease (T. whippeli), biopsy tissue from small bowel, abdominal or peripheral lymph nodes, or other organs of persons with signs and symptoms, to establish the diagnosis
  - ee. *Yersinia Pestis*

### **Policy Guidelines**

A discussion of every infectious agent that might be detected with a probe technique is beyond the scope of this policy. Many probes have been combined into panels of tests. For

the purposes of this policy, other than the respiratory virus panel, only individual probes are reviewed.

#### IV. Scientific Background

Nucleic acid hybridization technologies, including polymerase chain reaction (PCR), ligase- or helicase-dependent amplification, and transcription-mediated amplification, are beneficial tools for pathogen detection in blood culture and other clinical specimens due to high specificity and sensitivity (Khan,

2014). The use of nucleic acid-based methods to detect bacterial pathogens in a clinical laboratory setting offers “increased sensitivity and specificity over traditional microbiological techniques” due to its specificity, sensitivity, reduction in time, and high-throughput capability; however, “contamination potential, lack of standardization or validation for some assays, complex interpretation of results, and increased cost are possible limitations of these tests” (Mothershed & Whitney, 2006).

#### V. Guidelines and Recommendations

Specific guidelines for testing of many organisms listed within the policy coverage criteria is found in the updated **2018 Infectious Diseases Society of America (IDSA)** guidelines and recommendations titled, *A Guide to Utilization of the Microbiology Laboratory for Diagnosis of Infectious Diseases: 2018 Update* by the Infectious Diseases Society of America and the American Society for Microbiology (Miller et al., 2018). “This document is organized by body system, although many organisms are capable of causing disease in >1 body system. There may be a redundant mention of some organisms because of their propensity to infect multiple sites. One of the unique features of this document is its ability to assist clinicians who have specific suspicions regarding possible etiologic agents causing a specific type of disease. When the term “clinician” is used throughout the document, it also includes other licensed, advanced practice providers. Another unique feature is that in most chapters, there are targeted recommendations and precautions regarding selecting and collecting specimens for analysis for a disease process. It is very easy to access critical information about a specific body site just by consulting the table of contents. Within each chapter, there is a table describing the specimen needs regarding a variety of etiologic agents that one may suspect as causing the illness. The test methods in the tables are listed in priority order according to the recommendations of the authors and reviewers (Miller et al., 2018).”

##### Centers of Disease Control and Prevention (CDC)

###### *MRSA*

The CDC remarks that nucleic acid amplification tests (NAATs, such as PCR) “can be used for direct detection of *mecA*, the most common gene mediating oxacillin resistance in staphylococci”, but will not detect novel resistance mechanisms or uncommon phenotypes (CDC, 2019d).

###### *Candida Auris (C. auris)*

The CDC writes that “Molecular methods based on sequencing the D1-D2 region of the 28S rDNA or the Internal Transcribed Region (ITS) of rDNA also can identify *C. auris*.” The CDC further notes that various PCR methods have been developed for identifying *C. auris* (CDC, 2020).

### *Chlamydia Pneumoniae (C. pneumoniae)*

The CDC writes that RT-PCR is the “preferred” method of detecting a *C. pneumoniae* infection, with qPCR preferred for an acute infection. The CDC further notes that a positive culture should be confirmed by a second test, such as PCR (CDC, 2019a).

### *Ebola*

The CDC states that for diagnosis of Ebola, “there must be a combination of symptoms suggestive of EVD AND a possible exposure to EVD within 21 days before the onset of symptoms”. The CDC notes that PCR is one of the most common diagnostic methods (CDC, 2019b).

### *Salmonella*

The CDC writes that diagnosis requires detection of the *Salmonella* bacteria, be it through culture or a “culture-independent diagnostic test (CIDT)” (CDC, 2019f).

### *Giardia*

The CDC states that fecal immunoassays may be used for detection of *Giardia*, but that only molecular testing can identify *Giardia* subtypes (CDC, 2015b).

### *Non-Polio Enterovirus*

The CDC remarks that their laboratories “routinely” perform qualitative testing for enteroviruses, parechoviruses, and uncommon picornaviruses (CDC, 2018a).

### *Respiratory Syncytial Virus (RSV)*

The CDC writes that real-time reverse transcriptase-polymerase chain reaction (rRT-PCR) is the most commonly used diagnostic test, both for infants/younger children as well as older children (CDC, 2018b).

### *Mycoplasma Genitalium*

The CDC writes that “NAAT is the preferred method for *M. genitalium* detection” (CDC, 2015a).

### *Miscellaneous*

The CDC does not mention the need to quantify [through PCR] *Bartonella*, *Legionella pneumophila* or *Mycoplasma pneumoniae* (CDC, 2016, 2019c, 2019e). No guidance was found on Hepatitis G.

### **Committee on Infectious Diseases, American Academy of Pediatrics, 31st Edition (2018-2021, Red Book)**

The Committee on Infectious Diseases released joint guidelines with the American Academy of Pediatrics. In it, they note that “the presumptive diagnosis of mucocutaneous candidiasis or thrush usually can be made clinically”. They also state that FISH probes may rapidly detect *Candida* species from positive blood culture samples, although PCR assays have also been developed for this purpose (Pediatrics, 2018).

## VI. State and Federal Regulations, as applicable

As of 05/11/2020, a list of current U.S. Food and Drug Administration (FDA, 2020) approved or cleared nucleic acid-based microbial tests is available at: <https://www.fda.gov/medical-devices/vitrodiagnostics/nucleic-acid-based-tests>.

## VII. Applicable CPT/HCPCS Procedure Codes

*Billing applicable codes is not a guarantee of payment; see Section III for indications and limitations of coverage that may affect payment*

CPT	Code Description
87471	Infectious agent detection by nucleic acid (DNA or RNA); Bartonella henselae and Bartonella quintana, amplified probe technique
87472	Infectious agent detection by nucleic acid (DNA or RNA); Bartonella henselae and Bartonella quintana, quantification
87480	Infectious agent detection by nucleic acid (DNA or RNA); Candida species, direct probe technique
87481	Infectious agent detection by nucleic acid (DNA or RNA); Candida species, amplified probe technique
87482	Infectious agent detection by nucleic acid (DNA or RNA); Candida species, quantification
87485	Infectious agent detection by nucleic acid (DNA or RNA); Chlamydia pneumoniae, direct probe technique
87486	Infectious agent detection by nucleic acid (DNA or RNA); Chlamydia pneumoniae, amplified probe technique
87487	Infectious agent detection by nucleic acid (DNA or RNA); Chlamydia pneumoniae, quantification
87493	Infectious agent detection by nucleic acid (DNA or RNA); Clostridium difficile, toxin gene(s), amplified probe technique
87495	Infectious agent detection by nucleic acid (DNA or RNA); cytomegalovirus, direct probe technique

87496	Infectious agent detection by nucleic acid (DNA or RNA); cytomegalovirus, amplified probe technique
87497	Infectious agent detection by nucleic acid (DNA or RNA); cytomegalovirus, quantification
87498	Infectious agent detection by nucleic acid (DNA or RNA); enterovirus, amplified probe technique, includes reverse transcription when performed
87500	Infectious agent detection by nucleic acid (DNA or RNA); vancomycin resistance (eg, enterococcus species van A, van B), amplified probe technique
87516	Infectious agent detection by nucleic acid (DNA or RNA); hepatitis B virus, amplified probe technique
87517	Infectious agent detection by nucleic acid (DNA or RNA); hepatitis B virus, quantification
87525	Infectious agent detection by nucleic acid (DNA or RNA); hepatitis G, direct probe technique
87526	Infectious agent detection by nucleic acid (DNA or RNA); hepatitis G, amplified probe technique
87527	Infectious agent detection by nucleic acid (DNA or RNA); hepatitis G, quantification
87531	Infectious agent detection by nucleic acid (DNA or RNA); Herpes virus-6, direct probe technique
87532	Infectious agent detection by nucleic acid (DNA or RNA); Herpes virus-6, amplified probe technique
87533	Infectious agent detection by nucleic acid (DNA or RNA); Herpes virus-6, quantification
87540	Infectious agent detection by nucleic acid (DNA or RNA); Legionella pneumophila, direct probe technique
87541	Infectious agent detection by nucleic acid (DNA or RNA); Legionella pneumophila, amplified probe technique



87542	Infectious agent detection by nucleic acid (DNA or RNA); Legionella pneumophila, quantification
87563	Infectious agent detection by nucleic acid (DNA or RNA); Mycoplasma genitalium, amplified probe technique
87580	Infectious agent detection by nucleic acid (DNA or RNA); Mycoplasma pneumoniae, direct probe technique
87581	Infectious agent detection by nucleic acid (DNA or RNA); Mycoplasma pneumoniae, amplified probe technique
87582	Infectious agent detection by nucleic acid (DNA or RNA); Mycoplasma pneumoniae, quantification
87634	Infectious agent detection by nucleic acid (DNA or RNA); respiratory syncytial virus, amplified probe technique
87640	Infectious agent detection by nucleic acid (DNA or RNA); Staphylococcus aureus, amplified probe technique
87641	Infectious agent detection by nucleic acid (DNA or RNA); Staphylococcus aureus, methicillin resistant, amplified probe technique
87797	Infectious agent detection by nucleic acid (DNA or RNA), not otherwise specified; direct probe technique, each organism
87798	Infectious agent detection by nucleic acid (DNA or RNA), not otherwise specified; amplified probe technique, each organism
87799	Infectious agent detection by nucleic acid (DNA or RNA), not otherwise specified; quantification, each organism

Current Procedural Terminology© American Medical Association. All Rights reserved.

*Procedure codes appearing in Medical Policy documents are included only as a general reference tool for each policy. They may not be all-inclusive.*

## VIII. Evidence-based Scientific References

- CDC. (2015a). 2015 Sexually Transmitted Diseases Treatment Guidelines, Emerging Issues. Retrieved from <https://www.cdc.gov/std/tg2015/emerging.htm#myco>
- CDC. (2015b). Parasites - Giardia, Diagnosis & Detection. Retrieved from <https://www.cdc.gov/parasites/giardia/diagnosis.html>
- CDC. (2016). Bartonella Infection (Cat Scratch Disease, Trench Fever, and Carrion's Disease). Retrieved from <https://www.cdc.gov/bartonella/testing-faq/index.html>
- CDC. (2018a). Non-Polio Enterovirus, CDC Laboratory Testing & Procedures. Retrieved from <https://www.cdc.gov/non-polio-enterovirus/lab-testing/testing-procedures.html>
- CDC. (2018b). Respiratory Syncytial Virus Infection (RSV), For Healthcare Professionals. Retrieved from <https://www.cdc.gov/rsv/clinical/index.html#lab>
- CDC. (2019a). Chlamydia pneumoniae Infection, Diagnostic Methods. Retrieved from <https://www.cdc.gov/pneumonia/atypical/cpneumoniae/hcp/diagnostic.html>
- CDC. (2019b). Ebola (Ebola Virus Disease), Diagnosis. Retrieved from <https://www.cdc.gov/vhf/ebola/diagnosis/index.html>
- CDC. (2019c). Legionella (Legionnaires' Disease and Pontiac Fever). Retrieved from <https://www.cdc.gov/legionella/clinicians/diagnostic-testing.html>
- CDC. (2019d). Methicillin-resistant Staphylococcus aureus (MRSA), Laboratory Testing. Retrieved from [https://www.cdc.gov/mrsa/lab/index.html#anchor\\_1548439781](https://www.cdc.gov/mrsa/lab/index.html#anchor_1548439781)
- CDC. (2019e). Mycoplasma pneumoniae Infections. Retrieved from <https://www.cdc.gov/pneumonia/atypical/mycoplasma/hcp/diagnostic-methods.html>
- CDC. (2019f). Salmonella, Diagnostic and Public Health Testing. Retrieved from [https://www.cdc.gov/salmonella/general/diagnosis-treatment.html?CDC\\_AA\\_refVal=https%3A%2F%2Fwww.cdc.gov%2Fsalmonella%2Fgeneral%2Fdiagnosis.html](https://www.cdc.gov/salmonella/general/diagnosis-treatment.html?CDC_AA_refVal=https%3A%2F%2Fwww.cdc.gov%2Fsalmonella%2Fgeneral%2Fdiagnosis.html)
- CDC. (2020). Identification of Candida auris. Retrieved from [https://www.cdc.gov/fungal/candidaauris/identification.html?CDC\\_AA\\_refVal=https%3A%2F%2Fwww.cdc.gov%2Ffungal%2Fcandidaauris%2Frecommendations.html](https://www.cdc.gov/fungal/candidaauris/identification.html?CDC_AA_refVal=https%3A%2F%2Fwww.cdc.gov%2Ffungal%2Fcandidaauris%2Frecommendations.html)
- FDA. (2020). Nucleic Acid Based Tests. Retrieved from <https://www.fda.gov/medical-devices/vitrodiagnostics/nucleic-acid-based-tests>
- Khan, A. (2014). Rapid Advances in Nucleic Acid Technologies for Detection and Diagnostics of Pathogens. *J Microbiol Exp*, 1(2). doi:10.15406/jmen.2014.01.00009
- Miller, J. M., Binnicker, M. J., Campbell, S., Carroll, K. C., Chapin, K. C., Gilligan, P. H., . . . Yao, J. D. (2018). A Guide to Utilization of the Microbiology Laboratory for Diagnosis of Infectious Diseases: 2018 Update by the Infectious Diseases Society of America and the American Society for Microbiology. *Clinical Infectious Diseases*, ciy381-ciy381. doi:10.1093/cid/ciy381
- Mothershed, E. A., & Whitney, A. M. (2006). Nucleic acid-based methods for the detection of bacterial pathogens: present and future considerations for the clinical laboratory. *Clin Chim Acta*, 363(12), 206-220. doi:10.1016/j.cccn.2005.05.050
- Pediatrics, C. o. I. D. A. A. o. (2018). *Red Book*® 2018.

**IX. Revision History**

Revision Date	Summary of Changes
06-01-2021	Initial presentation