

Immunohistochemistry

Policy Number: AHS –P2018 – Immunohistochemistry	Prior Policy Name and Number, as applicable:
Effective Date: 06/01/2023	

[POLICY DESCRIPTION | INDICATIONS AND/OR LIMITATIONS OF COVERAGE | TABLE OF TERMINOLOGY | SCIENTIFIC BACKGROUND | GUIDELINES AND RECOMMENDATIONS | APPLICABLE STATE AND FEDERAL REGULATIONS | APPLICABLE CPT/HCPCS PROCEDURE CODES | EVIDENCE-BASED SCIENTIFIC REFERENCES | REVISION HISTORY](#)

I. Policy Description

Immunohistochemistry (IHC) is a very sensitive and specific staining technique that uses anatomical, biochemical, and immunological methods to identify cells, tissues, and organisms by the interaction of target antigens with highly specific monoclonal antibodies and visualization through the use of a biochemical tag or label (Fitzgibbons et al., 2014).

II. Indications and/or Limitations of Coverage

Application of coverage criteria is dependent upon an individual’s benefit coverage at the time of the request.

- 1) Code 88342 should be used for the first single antibody procedure and is reimbursed at one unit per specimen, up to four specimens, per date of service.
- 2) Code 88341 should be used for each additional single antibody per specimen and is reimbursed up to a maximum of 13 units per date of service.
- 3) Code 88344 should be used for each multiplex antibody per specimen, up to six specimens, per date of service.

III. Table of Terminology

Term	Definition
AFP	Alpha-fetoprotein
ASCO	The American Society of Clinical Oncology
<i>Bcl2</i>	<i>BCL2 apoptosis regulator</i>
b-HCG	Beta human chorionic gonadotropin
CAIX	Carbonic anhydrase IX
CAP	College of American Pathologists

Term	Definition
CD1a	Cluster of differentiation 1a
CD5	Cluster of differentiation 5
CD10	Cluster of differentiation 10
CD21	Cluster of differentiation 21
CD30	Cluster of differentiation 30
CD31	Cluster of differentiation 31
CD34	Cluster of differentiation 34
CD35	Cluster of differentiation 35
CD43	Cluster of differentiation 43
CD56	Cluster of differentiation 56
CD99	Cluster of differentiation 99
CD117	Cluster of differentiation 117
<i>CDH17</i>	<i>Cadherin-17</i>
<i>CDK4</i>	<i>Cyclin-dependent kinase 4</i>
CDX2	Caudal-type homeobox 2
CEA	Carcinoembryonic antigen
CK	Creatine kinase
CK17	Cytokeratin 17
CK20	Cytokeratin 20
CK5/6	Cytokeratin 5/6
CK903	Cytokeratin 903
CLIA'88	Clinical Laboratory Improvement Amendments of 1988
CMS	Centres for Medicare and Medicaid
CRC	Colorectal cancer
D2-40	Anti-Podoplanin
DNA	Deoxyribonucleic acid
DOG1	Delay of germination 1
ERG	ETS-related gene
ESMO	The European Society of Medical Oncology
FDA	Food and Drug Administration
FISH	Fluorescence in situ hybridization
Fli-1	Friend leukemia integration 1
FOXL2	Forkhead box protein L2
GATA3	GATA binding protein 3
GCDFP15	Gross cystic disease fluid protein 15
GI	Gastrointestinal tract
HepPar-1	General hepatocyte paraffin 1
<i>HER2</i>	<i>Human epidermal growth factor receptor 2</i>
HMB-45	Human melanoma black-45

Term	Definition
HNF-1b	Hepatocyte nuclear factor 1 beta
HPV	Human papillomavirus
IHC	Immunohistochemistry
IMP3	U3 small nucleolar ribonucleoprotein protein IMP3
INI1	Integrase interactor 1
ISH	In situ hybridization
KIM-1	Kidney injury molecule-1
LDTs	Laboratory developed tests
Maspin	Mammary serine protease inhibitor
MCPyV	Merkel cell polyomavirus
MDM2	Mouse double minute 2 homolog
MIB-1	MIB E3 ubiquitin protein ligase 1
mIHC	Multiplex immunohistochemistry
MiTF	Microphthalmia-associated transcription factor
MLH1	MutL homolog 1
MMR	Mismatch repair protein
MPO	Myeloperoxidase
MSA	Mammary serum antigen
MSH2	Mismatch repair protein Msh2
MSI	Microsatellite instability
MUC4	Mucin 4
MUC5AC	Mucin 5AC
MyoD1	Myoblast determination protein 1
<i>NANOG</i>	<i>Nanog Homeobox</i>
napsin A	Novel aspartic proteinase of the pepsin family A
NCCN	The National Cancer Coalition Network
NKX2.2	Homeobox protein
NKX3.1	Homeobox protein
NY-ESO-1	New York esophageal squamous cell carcinoma 1
OCT4	Octamer-binding transcription factor 4
p16	Cyclin-dependent kinase inhibitor 2A
p40	Protein subunit
P504S	Cytoplasmic protein
p63	Tumor protein p63
pan-Trk	Pan-tropomyosin-related-kinase
PAX2	Paired box 2
PAX8	Paired box 8
PDX1	Insulin promoter factor 1
PNET	Primitive neuro-ectodermal tumor

Term	Definition
PSA	Prostate-specific antigen
PSAP	Phosphoserine aminotransferase
PTEN	Phosphatase and tensin homolog
pVHL	Von hippel–lindau tumor suppressor
RB	Retinoblastoma protein
RCC	Renal cell carcinoma
RCCma	Renal cell carcinoma marker
S100P	S100 calcium-binding protein p
SALL4	Sal-like protein 4
SATB2	Special AT-rich sequence-binding protein 2
SF-1	Steroidogenic factor 1
SOX10	SRY-box transcription factor 10
TFE3	Transcription factor E3
TLE1	Transducin-like enhancer protein 1
TTF1	Transcription termination factor, RNA polymerase I
UPII	Uroplakin II
WT1	Wilms tumor protein

IV. Scientific Background

Immunohistochemistry (IHC) is used to identify certain components of tissues or cells (aka immunocytochemistry) via use of specific antibodies that can be visualized through a staining technique. The premise behind IHC is that distinct tissues and cells contain a unique set of antigens that allows them to be identified and differentiated. The selection of antibodies used for the evaluation of a specimen varies by the source of the specimen, the question to be answered, and the pathologist performing the test.

Importantly, an entirely sensitive and specific IHC marker rarely exists, and therefore, determinations are typically based on a pattern of positive and negative stains for a panel of several antibodies. The four most common IHC staining patterns include nuclear staining, cytoplasmic staining, membrane staining, and extracellular staining (Tuffaha et al., 2018). A single IHC marker approach (other than for pathogens such as cytomegalovirus or BK virus) is strongly discouraged since aberrant expression of a highly specific IHC marker can rarely occur. However, aberrant expression of the entire panel of highly specific IHC markers is nearly statistically impossible (Lin & Chen, 2014).

Multiplex immunohistochemistry (mIHC) is a particular IHC technique that allows multiple targets in a single tissue to be detected simultaneously; this approach is able to characterize “the tumor microenvironment including vascular architecture and hypoxia, cellular proliferation, cell death as well as drug distribution” (Kalra & Baker, 2017). Hence, mIHC can assist in the development of parameter tumor maps. Other researchers have utilized mIHC for its novel ability

to provide quantitative data on different types of tumor-infiltrating immune cells within a single tissue; this may improve cancer patient immunotherapy stratification (Hofman et al., 2019).

Clinical Utility and Validity

IHC can be used for a variety of purposes including: differentiation of benign from malignant tissue, differentiation among several types of cancer, selection of therapy, identification of the origin of a metastatic cancer, and identification of infectious organisms (Shah et al., 2012). IHC has many uses in the realm of tumor identification, and it has even been clinically used to pinpoint various breast cancer-specific markers, such as progesterone and estrogen receptors, gross cystic duct fluid protein, and mammaglobin (Hainsworth & Greco, 2022). Further, overexpression of the *HER2* oncogene, a predicative breast cancer biomarker, is often identified via IHC (Yamauchi & Hayes, 2022). In regards to tumor identification, a specific type of IHC, known as pan-Trk IHC, has been shown to positively identify inflammatory myofibroblastic tumors with a nuclear and cytoplasmic staining pattern that may assist in targeted therapy (Yamamoto et al., 2019).

Antibodies for use in IHC are available as single antibody reagents or in mixtures of a combination of antibodies. More than 200 diagnostic antibodies are generally available in a large clinical IHC laboratory, and hundreds of antibodies are usually available in research laboratories. The list of new antibodies is growing rapidly with the discovery of new biomarkers by molecular methodologies (Lizotte et al., 2016). Several studies have shown that a relatively low number of antibodies are capable of accurately diagnosing specific cancers and identifying the primary source of a metastasis (Le Stang et al., 2019; Lizotte et al., 2016; Prok & Prayson, 2006).

Common markers to identify tumor origin (Lin & Chen, 2014):

Primary Site	Markers
Lung adenocarcinoma	TTF1, napsin A
Breast carcinoma	GATA3, ER, GCDFP15
Urothelial carcinoma	GATA3, UPII, S100P, CK903, p63
Squamous cell carcinoma	p40, CK5/6
RCC, clear cell type	PAX8, RCCma, pVHL, KIM-1
Papillary RCC	P504S, RCCma, pVHL, PAX8, KIM-1
Translocational RCC	TFE3
Hepatocellular carcinoma	Arginase-1, glypican-3, HepPar-1
Adrenal cortical neoplasm	Mart-1, inhibin-a, calretinin, SF-1
Melanoma	S100, Mart-1, HMB-45, MiTF, SOX10
Merkel cell carcinoma	CK20 (perinuclear dot staining), MCPyV

Mesothelial origin	Calretinin, WT1, D2-40, CK5/6, mesothelin
Neuroendocrine origin	Chromogranin, synaptophysin, CD56
Upper GI tract	CDH17, CDX2, CK20
Lower GI tract	CDH17, SATB2, CDX2, CK20
Intrahepatic cholangiocarcinoma	pVHL, CAIX
Pancreas, acinar cell carcinoma	Glypican-3, antitrypsin
Pancreas, ductal adenocarcinoma	MUC5AC, CK17, Maspin, S100P, IMP3
Pancreas, neuroendocrine tumor	PR, PAX8, PDX1, CDH17, islet-1
Pancreas, solid pseudopapillary tumor	Nuclear b-catenin, loss of Ecadherin, PR, CD10, vimentin
Prostate, adenocarcinoma	PSA, NKX3.1, PSAP, ERG
Ovarian serous carcinoma	PAX8, ER, WT1
Ovarian clear cell carcinoma	pVHL, HNF-1b, KIM-1, PAX8
Endometrial stromal sarcoma	CD10, ER
Endometrial adenocarcinoma	PAX8/PAX2, ER, vimentin
Endocervical adenocarcinoma	PAX8, p16, CEA, HPV in situ hybridization, loss of PAX2
Thyroid follicular cell origin	TTF1, PAX8, thyroglobulin
Thyroid medullary carcinoma	Calcitonin, TTF1, CEA
Hyalinizing trabecular adenoma of the thyroid	MIB-1 (unique membranous staining pattern)
Salivary duct carcinoma	GATA3, AR, GCDFFP-15, HER2/neu
Thymic origin	PAX8, p63, CD5
Seminoma	SALL4, OCT4, CD117, D2-40
Yolk sac tumor	SALL4, glypican-3, AFP
Embryonal carcinoma	SALL4, OCT4, NANOG, CD30
Choriocarcinoma	b-HCG, CD10, SALL4
Sex cord–stromal tumors	SF-1, inhibin-a, calretinin, FOXL2
Vascular tumor	ERG, CD31, CD34, Fli-1
Synovial sarcoma	TLE1, cytokeratin

Chordoma	Cytokeratin, S100
Desmoplastic small round cell tumor	Cytokeratin, CD99, desmin, WT1 (N-terminus)
Alveolar soft part sarcoma	TFE3
Rhabdomyosarcoma	Myogenin, desmin, MyoD1
Smooth muscle tumor	SMA, MSA, desmin, calponin
Ewing sarcoma/PNET	NKX2.2, CD99, Fli-1
Myxoid and round cell liposarcoma	NY-ESO-1
Low-grade fibromyxoid sarcoma	MUC4
Epithelioid sarcoma	Loss of INI1, CD34, CK
Atypical lipomatous tumor	MDM2 (MDM2 by FISH is a more sensitive and specific test), CDK4
Histiocytosis X	CD1a, S100
Angiomyolipoma	HMB-45, SMA
Gastrointestinal stromal tumor	CD117, DOG1
Solitary fibrous tumor	CD34, Bcl2, CD99
Myoepithelial carcinoma	Cytokeratin and myoepithelial markers; may lose INI1
Myeloid sarcoma	CD43, CD34, MPO
Follicular dendritic cell tumor	CD21, CD35
Mast cell tumor	CD117, tryptase

V. Guidelines and Recommendations

Guidelines are lacking regarding the selection and number of antibodies that should be used for most immunohistochemistry evaluations. However, IHC is broadly used for conditions such as cancers, which are mentioned across many different societies. The below section is not a comprehensive list of guidance for immunohistochemistry.

College of American Pathologists (CAP)

CAP has published several reviews in Archives of Pathology & Laboratory Medicine that detail the quality control measures for IHC; further, CAP has also published more than 100 small IHC panels to address the frequently asked questions in diagnosis and differential diagnosis of specific

entities. These diagnostic panels are based on literature, IHC data, and personal experience. A single IHC marker approach (other than for pathogens such as cytomegalovirus or BK virus) is strongly discouraged since aberrant expression of a highly specific IHC marker can rarely occur. However, aberrant expression of the entire panel of highly specific IHC markers is nearly statistically impossible (Lin & Chen, 2014; Lin & Liu, 2014).

The American Society of Clinical Oncology (ASCO) and the College of American Pathologists (CAP)

ASCO and CAP currently recommend that “all newly diagnosed patients with breast cancer must have a HER2 test performed” (Wolff et al., 2013). Also, for those who develop metastatic disease, a HER2 test must be done on tissue from the metastatic site, if available. In less common HER2 breast cancer patterns, as observed in approximately 5% of cases by dual-probe in situ hybridization (ISH) assays, new recommendations have been made to make a final determination of positive or negative HER2 tissue. This new “diagnostic approach includes more rigorous interpretation criteria for ISH and requires concomitant IHC review for dual-probe ISH groups... to arrive at the most accurate HER2 status designation (positive or negative) based on combined interpretation of the ISH and IHC assays;” further, “The Expert Panel recommends that laboratories using single-probe ISH assays include concomitant IHC review as part of the interpretation of all single-probe ISH assay results” (Wolff et al., 2018).

The 2018 update included the following changes from the prior 2013 update, particularly focusing on infrequent HER2 test results that were of “uncertain biologic or clinical significance”:

- “Revision of the definition of IHC 2+ (equivocal) to the original FDA-approved criteria.
- Repeat HER2 testing on a surgical specimen if the initially tested core biopsy is negative is no longer stated as mandatory. A new HER2 test *may* (no longer *should*) be ordered on the excision specimen on the basis of some criteria (such as tumor grade 3).
- A more rigorous interpretation criteria of the less common patterns that can be seen in about 5% of all cases when HER2 status in breast cancer is evaluated using a dual-probe ISH testing. These cases, described as ISH groups 2 to 4, should now be assessed using a diagnostic approach that includes a concomitant review of the IHC test, which will help the pathologist make a final determination of the tumor specimen as HER2 positive or negative.

The Expert Panel also preferentially recommends the use of dual-probe instead of single-probe ISH assays, but it recognizes that several single-probe ISH assays have regulatory approval in many parts of the world” (Wolff et al., 2018)

The National Cancer Coalition Network

The NCCN has made numerous recommendations for use of IHC to diagnose and manage various types of cancer. Cancers with clinically useful IHC applications include breast, cervical, various leukemias, and colorectal cancer.

The NCCN states that the determination of estrogen receptor, progesterone receptor, and HER2 status for breast cancer is recommended and may be determined by IHC (NCCN, 2022a). Specifically, the NCCN guidelines state “consistent with the ASCO/CAP guidelines, the NCCN panel considers either IHC or ISH with either a single or dual probe as an acceptable method for making an initial determination of HER2 tumor status. PR testing by IHC on invasive cancers can aid in the prognostic classification of cancers and serve as a control for possible false negative ER results. Patients with ER-negative, PR-positive cancers may be considered for endocrine therapies, but the data on this group are noted to be limited” (NCCN, 2022a). Further, the NCCN recommendations concerning Lynch Syndrome (LS) state, “The panel recommends tumor testing with IHC and/or MSI be used as the primary approach for pathology-lab-based universal screening” (NCCN, 2022b). More recently, the NCCN has made additional recommendations to individuals diagnosed with any type of hereditary colorectal cancer (CRC) syndrome; these recommendations state that “all individuals newly diagnosed with CRC have either MSI or immunohistochemistry (IHC) testing for absence of 1 of the 4 DNA MMR proteins” (NCCN, 2022b).

The European Society of Medical Oncology (ESMO)

The ESMO recommends that for cancers of unknown primary, “immunohistochemistry should be applied meticulously in order to identify the tissue of origin and to exclude chemosensitive and potentially curable tumors” (Fizazi et al., 2015).

VI. Applicable State and Federal Regulations

Food and Drug Administration (FDA)

Many labs have developed specific tests that they must validate and perform in house. These laboratory-developed tests (LDTs) are regulated by the Centers for Medicare and Medicaid (CMS) as high-complexity tests under the Clinical Laboratory Improvement Amendments of 1988 (CLIA '88). LDTs are not approved or cleared by the U. S. Food and Drug Administration; however, FDA clearance or approval is not currently required for clinical use.

Recently, four clinical IHC biomarker assays (PTEN, RB, MLH1, and MSH2) have been validated for use as biomarkers in a nationwide clinical trial; these assays were then approved by the FDA as laboratory-developed tests to assist in the treatment selection of patients in clinical trials (Khoury et al., 2018). This shows that IHC assays are currently being utilized with molecular tests to assist in therapeutic decisions.

VII. Applicable CPT/HCPCS Procedure Codes

Procedure codes appearing in medical policy documents are only included as a general reference. This list may not be all inclusive and is subject to updates. In addition, codes listed are not a guarantee of payment.

CPT	Code Description
88341	Immunohistochemistry or immunocytochemistry, per specimen; each additional single antibody stain procedure
88342	Immunohistochemistry or immunocytochemistry, per spec; initial single antibody stain
88344	Immunohistochemistry or immunocytochemistry, per specimen; each multiplex antibody stain procedure

Current Procedural Terminology® American Medical Association. All Rights reserved.

VIII. Evidence-based Scientific References

- Fitzgibbons, P. L., Bradley, L. A., Fatheree, L. A., Alsabeh, R., Fulton, R. S., Goldsmith, J. D., Haas, T. S., Karabakhtsian, R. G., Loykasek, P. A., Marolt, M. J., Shen, S. S., Smith, A. T., & Swanson, P. E. (2014). Principles of analytic validation of immunohistochemical assays: Guideline from the College of American Pathologists Pathology and Laboratory Quality Center. *Arch Pathol Lab Med*, 138(11), 1432-1443. <https://doi.org/10.5858/arpa.2013-0610-CP>
- Fizazi, K., Greco, F. A., Pavlidis, N., Daugaard, G., Oien, K., & Pentheroudakis, G. (2015). Cancers of unknown primary site: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Ann Oncol*, 26 Suppl 5, v133-138. <https://doi.org/10.1093/annonc/mdv305>
- Hainsworth, J., & Greco, A. (2022, 01/20/2022). *Overview of the classification and management of cancers of unknown primary site*. <https://www.uptodate.com/contents/overview-of-the-classification-and-management-of-cancers-of-unknown-primary-site>
- Hofman, P., Badoual, C., Henderson, F., Berland, L., Hamila, M., Long-Mira, E., Lassalle, S., Roussel, H., Hofman, V., Tartour, E., & Ilie, M. (2019). Multiplexed Immunohistochemistry for Molecular and Immune Profiling in Lung Cancer-Just About Ready for Prime-Time? *Cancers (Basel)*, 11(3). <https://doi.org/10.3390/cancers11030283>
- Kalra, J., & Baker, J. (2017). Multiplex Immunohistochemistry for Mapping the Tumor Microenvironment. *Methods Mol Biol*, 1554, 237-251. https://doi.org/10.1007/978-1-4939-6759-9_17
- Khoury, J. D., Wang, W. L., Prieto, V. G., Medeiros, L. J., Kalhor, N., Hameed, M., Broaddus, R., & Hamilton, S. R. (2018). Validation of Immunohistochemical Assays for Integral Biomarkers in the NCI-MATCH EAY131 Clinical Trial. *Clin Cancer Res*, 24(3), 521-531. <https://doi.org/10.1158/1078-0432.Ccr-17-1597>
- Le Stang, N., Burke, L., Blaizot, G., Gibbs, A. R., Lebailly, P., Clin, B., Girard, N., & Galateau-Salle, F. (2019). Differential Diagnosis of Epithelioid Malignant Mesothelioma With Lung

- and Breast Pleural Metastasis: A Systematic Review Compared With a Standardized Panel of Antibodies-A New Proposal That May Influence Pathologic Practice. *Arch Pathol Lab Med*. <https://doi.org/10.5858/arpa.2018-0457-OA>
- Lin, F., & Chen, Z. (2014). Standardization of diagnostic immunohistochemistry: literature review and geisinger experience. *Arch Pathol Lab Med*, 138(12), 1564-1577. <https://doi.org/10.5858/arpa.2014-0074-RA>
- Lin, F., & Liu, H. (2014). Immunohistochemistry in undifferentiated neoplasm/tumor of uncertain origin. *Arch Pathol Lab Med*, 138(12), 1583-1610. <https://doi.org/10.5858/arpa.2014-0061-RA>
- Lizotte, P. H., Ivanova, E. V., Awad, M. M., Jones, R. E., Keogh, L., Liu, H., Dries, R., Almonte, C., Herter-Sprie, G. S., Santos, A., Feeney, N. B., Paweletz, C. P., Kulkarni, M. M., Bass, A. J., Rustgi, A. K., Yuan, G. C., Kufe, D. W., Janne, P. A., Hammerman, P. S., . . . Wong, K. K. (2016). Multiparametric profiling of non-small-cell lung cancers reveals distinct immunophenotypes. *JCI Insight*, 1(14), e89014. <https://doi.org/10.1172/jci.insight.89014>
- NCCN. (2022a). *Breast Cancer*. National Comprehensive Cancer Network. https://www.nccn.org/professionals/physician_gls/pdf/breast.pdf
- NCCN. (2022b). *Genetic/Familial High-Risk Assessment: Colorectal*. https://www.nccn.org/professionals/physician_gls/pdf/genetics_colon.pdf
- Prok, A., & Prayson, R. (2006). Thyroid transcription factor-1 staining is useful in identifying brain metastases of pulmonary origin. <https://www.sciencedirect.com/science/article/abs/pii/S109291340500119X>
- Shah, A. A., Frierson, H. F., & Cathro, H. P. (2012). Analysis of Immunohistochemical Stain Usage in Different Pathology Practice Settings. <https://doi.org/10.1309/AJCPAGVTCKDXKK0X>
- Tuffaha, M. S. A., Guski, H., & Kristiansen, G. (2018). *Immunohistochemistry in Tumor Diagnostics*. Springer, Cham. https://link.springer.com/chapter/10.1007/978-3-319-53577-7_1#citeas
- Wolff, A. C., Hammond, M. E., Hicks, D. G., Dowsett, M., McShane, L. M., Allison, K. H., Allred, D. C., Bartlett, J. M., Bilous, M., Fitzgibbons, P., Hanna, W., Jenkins, R. B., Mangu, P. B., Paik, S., Perez, E. A., Press, M. F., Spears, P. A., Vance, G. H., Viale, G., & Hayes, D. F. (2013). Recommendations for human epidermal growth factor receptor 2 testing in breast cancer: American Society of Clinical Oncology/College of American Pathologists clinical practice guideline update. *J Clin Oncol*, 31(31), 3997-4013. <https://doi.org/10.1200/jco.2013.50.9984>
- Wolff, A. C., Hammond, M. E. H., Allison, K. H., Harvey, B. E., Mangu, P. B., Bartlett, J. M. S., Bilous, M., Ellis, I. O., Fitzgibbons, P., Hanna, W., Jenkins, R. B., Press, M. F., Spears, P. A., Vance, G. H., Viale, G., McShane, L. M., & Dowsett, M. (2018). Human Epidermal Growth Factor Receptor 2 Testing in Breast Cancer: American Society of Clinical Oncology/College of American Pathologists Clinical Practice Guideline Focused Update. *J Clin Oncol*, 36(20), 2105-2122. <https://doi.org/10.1200/jco.2018.77.8738>

Yamamoto, H., Nozaki, Y., Kohashi, K., Kinoshita, I., & Oda, Y. (2019). Diagnostic utility of pan-Trk immunohistochemistry for inflammatory myofibroblastic tumors. *Histopathology*. <https://doi.org/10.1111/his.14010>

Yamauchi, H., & Hayes, D. (2022, 06/29/2022). *HER2 and predicting response to therapy in breast cancer*. <https://www.uptodate.com/contents/her2-and-predicting-response-to-therapy-in-breast-cancer>

IX. Revision History

Revision Date	Summary of Changes
01/01/2023	Initial Effective Date
04/04/2023	Updated the background, guidelines and recommendations, and evidence-based scientific references. Literature review did not necessitate any modifications to coverage criteria. Committee approved 4/4/2023